

SFRP1 Antibody

Rabbit mAb

Catalog # AP91263

Product Information

Application	WB, IHC, IF, ICC, IHF
Primary Accession	Q8N474
Reactivity	Human
Clonality	Monoclonal
Other Names	Frizzled related protein 1; FRP1; FrzA; SARP2; Secreted apoptosis related protein 2; Secreted frizzled related protein 1; SFRP1;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	35386

Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human SFRP1
Description	SFRP proteins appear to act as tumor suppressors, with loss of expression or function correlating with many invasive forms of cancer. Deletion of the corresponding SFRP1 gene and promoter hypermethylation leading to gene silencing has been reported in a number of cancers. Abnormal expression of SFRP1 and other Wnt signaling proteins is associated with some cases of retinitis pigmentosa.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	SFRP1
Synonyms	FRP, FRP1, SARP2
Function	Soluble frizzled-related proteins (sFRPS) function as modulators of Wnt signaling through direct interaction with Wnts. They have a role in regulating cell growth and differentiation in specific cell types. SFRP1 decreases intracellular beta-catenin levels (By similarity). Has antiproliferative effects on vascular cells, in vitro and in vivo, and can induce, in vivo, an angiogenic response. In vascular cell cycle, delays the G1 phase and entry into the S phase (By similarity). In kidney development, inhibits tubule formation and bud growth in metanephroi (By similarity). Inhibits WNT1/WNT4-mediated TCF- dependent transcription.


Cellular Location

Secreted. Note=Cell membrane or extracellular matrix-associated. Released by heparin-binding

Tissue Location

Widely expressed. Absent from lung, liver and peripheral blood leukocytes. Highest levels in heart and fetal kidney Also expressed in testis, ovary, fetal brain and lung, leiomyoma cells, myometrial cells and vascular smooth muscle cells. Expressed in foreskin fibroblasts and in keratinocytes

Images



Western blot analysis of SFRP1 expression in A375 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.