

GATA3 Antibody

Rabbit mAb

Catalog # AP91270

Product Information

Application	WB, IHC, IF, FC, ICC, IP, IHF
Primary Accession	P23771
Reactivity	Human
Clonality	Monoclonal
Other Names	GATA-binding factor 3; GATA3;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	47916


Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 FC 1:100 ChIP
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human GATA3
Description	Transcriptional activator which binds to the enhancer of the T-cell receptor alpha and delta genes. Binds to the consensus sequence 5'-AGATAG-3'.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	GATA3
Function	Transcriptional activator which binds to the enhancer of the T-cell receptor alpha and delta genes. Binds to the consensus sequence 5'-AGATAG-3'. Required for the T-helper 2 (Th2) differentiation process following immune and inflammatory responses. Positively regulates ASB2 expression (By similarity). Coordinates macrophage transcriptional activation and UCP2-dependent metabolic reprogramming in response to IL33. Upon tissue injury, acts downstream of IL33 signaling to drive differentiation of inflammation-resolving alternatively activated macrophages.
Cellular Location	Nucleus.
Tissue Location	T-cells and endothelial cells.

Images



Western blot analysis of GATA3 expression in (1) SH-SY5Y cell lysate; (2) Jurkat cell lysate.

Image not found : 202311/AP91270-IHC.jpg

Immunohistochemical analysis of paraffin-embedded human bladder cancer, using GATA3 Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.