

G6PD Antibody

Rabbit mAb

Catalog # AP91271

Product Information

| | |
|--------------------------|--|
| Application | WB, IHC, IF, FC, ICC, IHF |
| Primary Accession | P11413 |
| Reactivity | Human |
| Clonality | Monoclonal |
| Other Names | G6PD; G6PD1; G6pdx; Glucose 6 phosphate 1 dehydrogenase; Glucose 6 phosphate dehydrogenase; Glucose 6 phosphate dehydrogenase, G6PD; MET19; POS10; Zwfp1p; |
| Isotype | Rabbit IgG |
| Host | Rabbit |
| Calculated MW | 59257 |


Additional Information

| | |
|-------------------------------------|--|
| Dilution | WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 FC 1:100 |
| Purification | Affinity-chromatography |
| Immunogen | A synthesized peptide derived from human G6PD |
| Description | Catalyzes the rate-limiting step of the oxidative pentose-phosphate pathway, which represents a route for the dissimilation of carbohydrates besides glycolysis. The main function of this enzyme is to provide reducing power (NADPH) and pentose phosphates for fatty acid and nucleic acid synthesis. |
| Storage Condition and Buffer | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |

Protein Information

| | |
|--------------------------|---|
| Name | G6PD |
| Function | Catalyzes the rate-limiting step of the oxidative pentose- phosphate pathway, which represents a route for the dissimilation of carbohydrates besides glycolysis. The main function of this enzyme is to provide reducing power (NADPH) and pentose phosphates for fatty acid and nucleic acid synthesis. |
| Cellular Location | Cytoplasm, cytosol. Membrane; Peripheral membrane protein |
| Tissue Location | Isoform Long is found in lymphoblasts, granulocytes and sperm |

Images



Western blot analysis of G6PD expression in MCF7 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.