10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Drosha Antibody

Rabbit mAb
Catalog # AP91300

Product Information

Application	WB, IHC, IF, FC, ICC, IHF
Primary Accession	Q9NRR4
Reactivity	Human
Clonality	Monoclonal
Other Names	Ribonuclease 3 (EC:3.1.26.3); Protein Drosha; Ribonuclease III; RNase III; p241; DROSHA; RN3; RNASE3L; RNASEN;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	159316

Additional Information

Dilution	WB 1:1000~1:5000 IHC 1:50~1:200 ICC/IF 1:50~1:200 FC 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Drosha
Description	Drosha was identified as a nuclear RNase III that catalyzes the initial step of microRNA (miRNA) processing. This enzyme processes the long primary transcript pri-miRNAs into stem-looped pre-miRNAs. Interference of Drosha results in the increase of pri-miRNAs and the decrease of pre-miRNAs. Drosha exists in a multiprotein complex called Microprocessor along with other components such as DGCR8. Drosha, along with DGCR8, is necessary for miRNA biogenesis.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information


Name	DROSHA
Synonyms	RN3, RNASE3L, RNASEN
Function	Ribonuclease III double-stranded (ds) RNA-specific endoribonuclease that is involved in the initial step of microRNA (miRNA) biogenesis. Component of the microprocessor complex that is required to process primary miRNA transcripts (pri-miRNAs) to release precursor miRNA (pre-miRNA) in the nucleus. Within the microprocessor complex, DROSHA cleaves the 3' and 5' strands of a stem-loop in pri- miRNAs (processing center 11 bp from the dsRNA-ssRNA junction) to release hairpin-shaped pre-miRNAs that are subsequently cut by the cytoplasmic DICER to generate mature miRNAs. Involved also in pre-rRNA processing. Cleaves double-strand RNA and does

not cleave single-strand RNA. Involved in the formation of GW bodies. Plays a role in growth homeostasis in response to autophagy in motor neurons (By similarity).

Cellular Location Nucleus. Nucleus, nucleolus. Cytoplasm {ECO:0000250|UniProtKB:Q5HZJ0}. Note=A fraction is translocated to the nucleolus during the S phase of the cell cycle. Localized in GW bodies (GWBs), also known as P-bodies.

Tissue Location Ubiquitous..

Images



Western blot analysis of Drosha expression in 293 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.