

DR5 Antibody

Rabbit mAb

Catalog # AP91307

Product Information

Application	WB, FC, IP
Primary Accession	O14763
Reactivity	Human
Clonality	Monoclonal
Other Names	Death receptor 5; TRAIL receptor 2; TRAIL-R2; CD262; TNFRSF10B; DR5; KILLER; TRAILR2; TRICK2; ZTNFR9;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	47878

Additional Information


Dilution	WB 1:1000~1:5000 IP 1:30 FC 1:100
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human DR5
Description	Plays an important role in the regulation of apoptosis in various physiological systems. Promotes the activation of NF-kappa-B. Essential for ER stress-induced apoptosis.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	TNFRSF10B
Synonyms	DR5, KILLER, TRAILR2, TRICK2, ZTNFR9
Function	Receptor for the cytotoxic ligand TNFSF10/TRAIL (PubMed: 10549288). The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. Promotes the activation of NF-kappa-B. Essential for ER stress-induced apoptosis.
Cellular Location	Membrane; Single-pass type I membrane protein.
Tissue Location	Widely expressed in adult and fetal tissues; very highly expressed in tumor cell lines such as HeLaS3, K-562, HL-60, SW480, A-549 and G-361; highly expressed in heart, peripheral blood lymphocytes, liver, pancreas, spleen, thymus, prostate, ovary, uterus, placenta, testis, esophagus, stomach and

throughout the intestinal tract; not detectable in brain

Images



Western blot analysis of DR5 expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.