10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Glypican 3 Antibody

Rabbit mAb
Catalog # AP91348

## Product Information

| | |
|---|---|
| **Application** | WB, FC |
| **Primary Accession** | P51654 |
| **Reactivity** | Human |
| **Clonality** | Monoclonal |
| **Other Names** | DGSX; Glypican proteoglycan 3; Gpc3; GTR2 2; MXR7; OCI5; SDYS; Secreted glypican-3; SGB; SGBS1; |
| **Isotype** | Rabbit IgG |
| **Host** | Rabbit |
| **Calculated MW** | 65563 |

## Additional Information


| | |
|---|---|
| **Dilution** | WB 1:500~1:2000 FC 1:50 |
| **Purification** | Affinity-chromatography |
| **Immunogen** | A synthesized peptide derived from human Glypican 3 |
| **Description** | Cell surface proteoglycan that bears heparan sulfate. Inhibits the dipeptidyl peptidase activity of DPP4. May be involved in the suppression/modulation of growth in the predominantly mesodermal tissues and organs. |
| **Storage Condition and Buffer** | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |

## Protein Information

| | |
|---|---|
| **Name** | GPC3 |
| **Synonyms** | OCI5 |
| **Function** | Cell surface proteoglycan (PubMed: 14610063). Negatively regulates the hedgehog signaling pathway when attached via the GPI- anchor to the cell surface by competing with the hedgehog receptor PTC1 for binding to hedgehog proteins (By similarity). Binding to the hedgehog protein SHH triggers internalization of the complex by endocytosis and its subsequent lysosomal degradation (By similarity). Positively regulates the canonical Wnt signaling pathway by binding to the Wnt receptor Frizzled and stimulating the binding of the Frizzled receptor to Wnt ligands (PubMed:16227623, PubMed:24496449). Positively regulates the non-canonical Wnt signaling pathway (By similarity). Binds to CD81 which decreases the availability of free CD81 for binding to the transcriptional repressor HHEX, resulting in nuclear translocation of HHEX and transcriptional repression (By similarity). Inhibits the dipeptidyl peptidase activity of DPP4 (PubMed:17549790). Plays a role in |

| | |
|---|---|
| | limb patterning and skeletal development by controlling the cellular response to BMP4 (By similarity). Modulates the effects of growth factors BMP2, BMP7 and FGF7 on renal branching morphogenesis (By similarity). Required for coronary vascular development (By similarity). Plays a role in regulating cell movements during gastrulation (By similarity). |
| **Cellular Location** | Cell membrane; Lipid-anchor, GPI-anchor {ECO:0000250\|UniProtKB:P13265}; Extracellular side {ECO:0000250\|UniProtKB:P13265} |
| **Tissue Location** | Detected in placenta (at protein level) (PubMed:32337544). Highly expressed in lung, liver and kidney |

## Images




Western blot analysis of Glypican 3 expression in HepG2 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.