

FTMT Antibody

Rabbit mAb

Catalog # AP91391

Product Information

Application	WB
Primary Accession	Q8N4E7
Reactivity	Human
Clonality	Monoclonal
Other Names	FerH; Ferritin; Ferritin H subunit; Ferritin, mitochondrial; Fth3; FTMT; mitochondrial; Mitochondrial ferritin; MTF;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	27538


Additional Information

Dilution	WB 1:500~1:2000
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human FTMT
Description	Stores iron in a soluble, non-toxic, readily available form. Important for iron homeostasis. Has ferroxidase activity. Iron is taken up in the ferrous form and deposited as ferric hydroxides after oxidation.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	FTMT (HGNC:17345)
Function	Catalyzes the oxidation of ferrous iron(II) to ferric iron(III) and stores iron in a soluble, non-toxic, readily available form (PubMed: 11323407 , PubMed: 15201052). Important for iron homeostasis (PubMed: 11323407 , PubMed: 15201052). Iron is taken up in the ferrous form and deposited as ferric hydroxides after oxidation (PubMed: 11323407 , PubMed: 15201052).
Cellular Location	Mitochondrion.
Tissue Location	Detected in testis and erythroleukemia. Expression is very low or not detectable in brain, colon, heart, kidney, liver, lung, muscle, placental, spleen and small intestine

Images



Western blot analysis of FTMT expression in HepG2 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.