

OGG1 Antibody

Rabbit mAb

Catalog # AP91396

Product Information

Application	WB
Primary Accession	O15527
Reactivity	Human
Clonality	Monoclonal
Other Names	8 hydroxyguanine DNA glycosylase; AP lyase; HMMH; HOGG1; MMH; MUTM; N glycosylase; Ogg1; OGH 1;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	38782


Additional Information

Dilution	WB 1:500~1:2000
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human OGG1
Description	DNA repair enzyme that incises DNA at 8-oxoG residues. Excises 7,8-dihydro-8-oxoguanine and 2,6-diamino-4-hydroxy-5-N-methylformamidopyrimidine (FAPY) from damaged DNA. Has a beta-lyase activity that nicks DNA 3' to the lesion.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	OGG1
Synonyms	MMH, MUTM, OGH1
Function	DNA repair enzyme that incises DNA at 8-oxoG residues. Excises 7,8-dihydro-8-oxoguanine and 2,6-diamino-4-hydroxy-5-N-methylformamidopyrimidine (FAPY) from damaged DNA. Has a beta-lyase activity that nicks DNA 3' to the lesion.
Cellular Location	Nucleus, nucleoplasm. Nucleus speckle. Nucleus matrix. Note=Together with APEX1 is recruited to nuclear speckles in UVA-irradiated cells [Isoform 2A]; Mitochondrion.
Tissue Location	Ubiquitous.

Images



Western blot analysis of OGG1 expression in HepG2 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.