

TXNRD1 Antibody

Rabbit mAb Catalog # AP91399

Product Information

Application Primary Accession Reactivity Clonality Other Names	WB, IHC, IF, FC, ICC, IP, IHF <u>Q16881</u> Rat, Human, Mouse Monoclonal GRIM12; KDRF; KM 102 derived reductase like factor; xidoreductase; Thioredoxin reductase 1; TR1; TRXR1; TXNR; TXNRD1
lsotype	Rabbit IgG
Host	Rabbit
Calculated MW	70906

Additional Information

Dilution Purification Immunogen Description	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 IP 1:50 FC 1:120 Affinity-chromatography A synthesized peptide derived from human TXNRD1 Isoform 1 may possess glutaredoxin activity as well as thioredoxin reductase activity and induces actin and tubulin polymerization, leading to formation of cell membrane protrusions. Isoform 4 enhances the transcriptional activity of estrogen receptors alpha and beta while isoform 5 enhances the transcriptional activity of the beta receptor only. Isoform 5 also mediates cell death induced by a combination of interferon-beta and retinoic acid.
Storage Condition and Buffer	

Protein Information

Name	TXNRD1 (<u>HGNC:12437</u>)
Synonyms	GRIM12, KDRF
Function	Reduces disulfideprotein thioredoxin (Trx) to its dithiol- containing form (PubMed: <u>8577704</u>). Homodimeric flavoprotein involved in the regulation of cellular redox reactions, growth and differentiation. Contains a selenocysteine residue at the C-terminal active site that is essential for catalysis (Probable). Also has reductase activity on hydrogen peroxide (H2O2) (PubMed: <u>10849437</u>).
Cellular Location	[Isoform 1]: Cytoplasm [Isoform 5]: Cytoplasm
Tissue Location	[Isoform 1]: Expressed predominantly in Leydig cells (at protein level). Also expressed in ovary, spleen, heart, liver, kidney and pancreas and in a number





Western blot analysis of TXNRD1 expression in (1) Jurkat cell lysate; (2) NIH/3T3 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.