

# **DDR2** Antibody

Rabbit mAb Catalog # AP91424

### **Product Information**

Application WB Primary Accession Q16832

Reactivity Rat, Human, Mouse

**Clonality** Monoclonal

Other Names CD167b; NTRKR3; Receptor related 3; TKT;

IsotypeRabbit IgGHostRabbitCalculated MW96736

## **Additional Information**

**Dilution** WB 1:500~1:2000

**Purification** Affinity-chromatography

**Immunogen** A synthesized peptide derived from human DDR2

**Description**This tyrosine kinase receptor for fibrillar collagen mediates fibroblast

migration and proliferation. Contributes to cutaneous wound healing.

**Storage Condition and Buffer** Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

#### **Protein Information**

Name DDR2

Synonyms NTRKR3, TKT, TYRO10

**Function** Tyrosine kinase involved in the regulation of tissues remodeling

(PubMed:30449416). It functions as a cell surface receptor for fibrillar collagen and regulates cell differentiation, remodeling of the extracellular matrix, cell migration and cell proliferation. Required for normal bone development. Regulates osteoblast differentiation and chondrocyte

maturation via a signaling pathway that involves MAP kinases and leads to the activation of the transcription factor RUNX2. Regulates remodeling of the extracellular matrix by up- regulation of the collagenases MMP1, MMP2 and MMP13, and thereby facilitates cell migration and tumor cell invasion. Promotes fibroblast migration and proliferation, and thereby contributes to


cutaneous wound healing.

**Cellular Location** Cell membrane; Single-pass type I membrane protein

**Tissue Location** Detected in osteocytes, osteoblastic cells in subchondral bone, bone lining

cells, tibia and cartilage (at protein level). Detected at high levels in heart and lung, and at low levels in brain, placenta, liver, skeletal muscle, pancreas, and kidney

# **Images**



Western blot analysis of DDR2 expression in (1) Jurkat cell lysate; (2) NIH/3T3 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.