

mSin3A Antibody

Rabbit mAb

Catalog # AP91428

Product Information

Application	WB, IHC, IF, FC, ICC, IHF
Primary Accession	Q96ST3
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	Histone deacetylase complex subunit Sin 3a; Paired amphipathic helix protein Sin 3a; SIN3 homolog A;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	145175

Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 FC 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human mSin3A
Description	Acts as a transcriptional repressor. Interacts with MXI1 to repress MYC responsive genes and antagonize MYC oncogenic activities. Also interacts with MAD-MAX heterodimers by binding to MAD. The heterodimer then represses transcription by tethering SIN3A to DNA. Acts as a corepressor for REST.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information


Name	SIN3A (HGNC:19353)
Function	Acts as a transcriptional repressor. Corepressor for REST. Interacts with MXI1 to repress MYC responsive genes and antagonize MYC oncogenic activities. Also interacts with MXD1-MAX heterodimers to repress transcription by tethering SIN3A to DNA. Acts cooperatively with OGT to repress transcription in parallel with histone deacetylation. Involved in the control of the circadian rhythms. Required for the transcriptional repression of circadian target genes, such as PER1, mediated by the large PER complex through histone deacetylation. Cooperates with FOXK1 to regulate cell cycle progression probably by repressing cell cycle inhibitor genes expression (By similarity). Required for cortical neuron differentiation and callosal axon elongation (By similarity).
Cellular Location	Nucleus {ECO:0000255 PROSITE-ProRule:PRU00810, ECO:0000269 PubMed:16820529}. Nucleus, nucleolus. Note=Recruited to the

nucleolus by SAP30L

Tissue Location

Expressed in the developing brain, with highest levels of expression detected in the ventricular zone of various cortical regions.

Images



Western blot analysis of mSin3A expression in (1) K562 cell lysate; (2) RAW 264.7 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.