

Angiopoietin 2 Antibody

Rabbit mAb

Catalog # AP91455

Product Information

Application	WB, IHC
Primary Accession	O15123
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	AGPT 2; Agpt2; ANG2; Angiopoietin 2a; Angiopoietin 2B; Angiopoietin2; ANGPT 2; Angpt2; Tie2 ligand;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	56919


Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Angiopoietin 2
Description	Can induce tyrosine phosphorylation of TIE2. Binds to TIE2 receptor and counteracts blood vessel maturation/stability mediated by angiopoietin-1. Its function may be context-dependent.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	ANGPT2
Function	Binds to TEK/TIE2, competing for the ANGPT1 binding site, and modulating ANGPT1 signaling (PubMed: 15284220 , PubMed: 19116766 , PubMed: 19223473 , PubMed: 9204896). Can induce tyrosine phosphorylation of TEK/TIE2 in the absence of ANGPT1 (PubMed: 15284220 , PubMed: 19116766 , PubMed: 19223473 , PubMed: 9204896). In the absence of angiogenic inducers, such as VEGF, ANGPT2-mediated loosening of cell-matrix contacts may induce endothelial cell apoptosis with consequent vascular regression. In concert with VEGF, it may facilitate endothelial cell migration and proliferation, thus serving as a permissive angiogenic signal (PubMed: 15284220 , PubMed: 19116766 , PubMed: 19223473 , PubMed: 9204896). Involved in the regulation of lymphangiogenesis (PubMed: 32908006).
Cellular Location	Secreted.

Images



Western blot analysis of Angiopoietin 2 expression in HUVEC cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.