

Cyclophilin F Antibody

Rabbit mAb Catalog # AP91459

Product Information

Application Primary Accession Reactivity Clonality Other Names	WB, IHC, IP <u>P30405</u> Rat, Human, Mouse Monoclonal Cyclophilin 3; cyclophilin D; Cyclophilin F; Cyp D; CyP M; CYP3; CypD; hCyP3; Mitochondrial cyclophilin; PPIase; PPIase F; Ppif; Rotamase;
lsotype	Rabbit IgG
Host	Rabbit
Calculated MW	22040

Additional Information

Dilution Purification Immunogen Description	WB 1:500~1:2000 IHC 1:50~1:200 IP 1:50 Affinity-chromatography A synthesized peptide derived from human Cyclophilin F PPIases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides.
Storage Condition and Buffer	

Protein Information

Name	PPIF
Synonyms	CYP3
Function	PPIase that catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and may therefore assist protein folding (PubMed:20676357). Involved in regulation of the mitochondrial permeability transition pore (mPTP) (PubMed:26387735). It is proposed that its association with the mPTP is masking a binding site for inhibiting inorganic phosphate (Pi) and promotes the open probability of the mPTP leading to apoptosis or necrosis; the requirement of the PPIase activity for this function is debated (PubMed:26387735). In cooperation with mitochondrial p53/TP53 is involved in activating oxidative stress-induced necrosis (PubMed:22726440). Involved in modulation of mitochondrial membrane F(1)F(0) ATP synthase activity and regulation of mitochondrial matrix adenine nucleotide levels (By similarity). Has anti-apoptotic activity independently of mPTP and in cooperation with BCL2 inhibits cytochrome c-dependent apoptosis (PubMed:19228691).

Images



Western blot analysis of Cyclophilin F expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.