

HSD11B1 Antibody

Rabbit mAb

Catalog # AP91472

Product Information

Application	WB, IHC
Primary Accession	P28845
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	11-DH; 11-beta-HSD1; HSD11B1; HSD11; HSD11L; 11 beta HSD1;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	32401

Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human HSD11B1
Description	Catalyzes reversibly the conversion of cortisol to the inactive metabolite cortisone. Catalyzes reversibly the conversion of 7-ketocholesterol to 7-beta-hydroxycholesterol. In intact cells, the reaction runs only in one direction, from 7-ketocholesterol to 7-beta-hydroxycholesterol.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	HSD11B1 (HGNC:5208)
Synonyms	HSD11, HSD11L, SDR26C1
Function	Controls the reversible conversion of biologically active glucocorticoids such as cortisone to cortisol, and 11- dehydrocorticosterone to corticosterone in the presence of NADP(H) (PubMed: 10497248 , PubMed: 12460758 , PubMed: 14973125 , PubMed: 15152005 , PubMed: 15280030 , PubMed: 17593962 , PubMed: 21453287 , PubMed: 27927697 , PubMed: 30902677). Participates in the corticosteroid receptor-mediated anti-inflammatory response, as well as metabolic and homeostatic processes (PubMed: 10497248 , PubMed: 12414862 , PubMed: 15152005 , PubMed: 21453287). Plays a role in the secretion of aqueous humor in the eye, maintaining a normotensive, intraocular environment (PubMed: 11481269). Bidirectional in vitro, predominantly functions as a reductase in vivo, thereby increasing the concentration of active glucocorticoids (PubMed: 10497248 , PubMed: 11481269 , PubMed: 12414862 , PubMed: 12460758). It has broad

substrate specificity, besides glucocorticoids, it accepts other steroid and sterol substrates (PubMed:[15095019](#), PubMed:[15152005](#), PubMed:[17593962](#), PubMed:[21453287](#)). Interconverts 7-oxo- and 7-hydroxy-neurosteroids such as 7- oxopregnenolone and 7beta-hydroxypregnenolone, 7-oxodehydroepiandrosterone (3beta-hydroxy-5-androstene-7,17-dione) and 7beta-hydroxydehydroepiandrosterone (3beta,7beta-dihydroxyandrost-5-en-17-one), among others (PubMed:[17593962](#)). Catalyzes the stereo-specific conversion of the major dietary oxysterol, 7-ketocholesterol (7-oxocholesterol), into the more polar 7-beta-hydroxycholesterol metabolite (PubMed:[15095019](#), PubMed:[15152005](#)). 7-oxocholesterol is one of the most important oxysterols, it participates in several events such as induction of apoptosis, accumulation in atherosclerotic lesions, lipid peroxidation, and induction of foam cell formation (PubMed:[15095019](#)). Mediates the 7-oxo reduction of 7-oxolithocholate mainly to chenodeoxycholate, and to a lesser extent to ursodeoxycholate, both in its free form and when conjugated to glycine or taurine, providing a link between glucocorticoid activation and bile acid metabolism (PubMed:[21453287](#)). Catalyzes the synthesis of 7-beta-25-dihydroxycholesterol from 7-oxo-25-hydroxycholesterol in vitro, which acts as a ligand for the G-protein-coupled receptor (GPCR) Epstein-Barr virus-induced gene 2 (EBI2) and may thereby regulate immune cell migration (PubMed:[30902677](#)).


Cellular Location

Endoplasmic reticulum membrane; Single-pass type II membrane protein

Tissue Location

Widely expressed, highest expression in liver, lower in testis, ovary, lung, foreskin fibroblasts, and much lower in kidney (PubMed:[1885595](#)). Expressed in liver (at protein level) (PubMed:[21453287](#)). Expressed in the basal cells of the corneal epithelium and in the ciliary nonpigmented epithelium (both at mRNA and at protein level) (PubMed:[11481269](#)).

Images



Western blot analysis of HSD11B1 expression in human fetal liver lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.