10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ULK3 Antibody

Rabbit mAb
Catalog # AP91482

Product Information

Application	WB, IHC, IF, FC, ICC, IHF
Primary Accession	Q6PHR2
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	Serine/threonine-protein kinase ULK3; Ulk3; unc 51 like kinase 3 (C. elegans); Unc-51-like kinase 3;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	53444

Additional Information


Dilution	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 FC 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human ULK3
Description	Serine/threonine protein kinase which enhances GLI1 and GLI2 transcriptional activity and consequently positively regulates GLI-dependent SHH signaling. May exert this function by promoting GLI1 nuclear localization. Phosphorylates in vitro GLI2, as well as GLI1 and GLI3, although less efficiently.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	ULK3 {ECO:0000303|PubMed:19878745, ECO:0000312|HGNC:HGNC:19703}
Function	Serine/threonine protein kinase that acts as a regulator of smoothened signaling and autophagy (PubMed:19279323, PubMed:19878745, PubMed:20643644, PubMed:31279575). Acts as a positive regulator of smoothened signaling in the presence of hedgehog (DHH, IHH or SHH) ligand: autophosphorylates and mediates phosphorylation of full-length GLI2, promoting subsequent phosphorylation by CDK1 and GLI2 dissociation from SUFU and translocation to the nucleus (PubMed:19878745, PubMed:20643644, PubMed:31279575). Also phosphorylates as GLI1 and GLI3, although less efficiently (PubMed:19878745, PubMed:20643644). In the absence of hedgehog ligand, interacts with SUFU, thereby inactivating the protein kinase activity and preventing phosphorylation of GLI proteins (GLI1, GLI2 and/or GLI3) (PubMed:20643644). Also acts as a regulator of autophagy: following cellular senescence, able to induce autophagy (PubMed:19279323).

Cellular Location	Cytoplasm. Note=Localizes to pre-autophagosomal structure during cellular senescence.
Tissue Location	Widely expressed. Highest levels observed in fetal brain. In adult tissues, high levels in brain, liver and kidney, moderate levels in testis and adrenal gland and low levels in heart, lung, stomach, thymus, prostate and placenta. In the brain, highest expression in the hippocampus, high levels also detected in the cerebellum, olfactory bulb and optic nerve. In the central nervous system, lowest levels in the spinal cord

Images



Western blot analysis of ULK3 expression in 293T cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.