

MSH2 Antibody

Rabbit mAb Catalog # AP91499

Product Information

Application Primary Accession Reactivity Clonality Other Names	WB, FC <u>P43246</u> Human Monoclonal DNA mismatch repair protein Msh2; hMSH2; MutS protein homolog 2; MSH2; COCA1; LCFS2;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	104743

Additional Information

Dilution Purification	WB 1:500~1:2000 FC 1:50 Affinity-chromatography
Immunogen	A synthesized peptide derived from human MSH2
Description	MSH2 (MutS homologue 2) forms the hMutS-α dimer with MSH6 and is an essential component of the mismatch repair process. hMutS-α is part of the BRCA1-associated surveillance complex (BASC), a complex that also contains BRCA1, MLH1, ATM, BLM, PMS2 proteins and the Rad50-Mre11-NBS1 complex.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name MSH2 Function Component of the post-replicative DNA mismatch repair system (MMR). Forms two different heterodimers: MutS alpha (MSH2-MSH6 heterodimer) and MutS beta (MSH2-MSH3 heterodimer) which binds to DNA mismatches thereby initiating DNA repair. When bound, heterodimers bend the DNA helix and shields approximately 20 base pairs. MutS alpha recognizes single base mismatches and dinucleotide insertion-deletion loops (IDL) in the DNA. MutS beta recognizes larger insertion-deletion loops up to 13 nucleotides long. After mismatch binding, MutS alpha or beta forms a ternary complex with the MutL alpha heterodimer, which is thought to be responsible for directing the downstream MMR events, including strand discrimination, excision, and resynthesis. Recruits DNA helicase MCM9 to chromatin which unwinds the mismatch containing DNA strand (PubMed: 26300262). ATP binding and hydrolysis play a pivotal role in mismatch repair functions. The ATPase activity

	associated with MutS alpha regulates binding similar to a molecular switch: mismatched DNA provokes ADP>ATP exchange, resulting in a discernible conformational transition that converts MutS alpha into a sliding clamp capable of hydrolysis-independent diffusion along the DNA backbone. This transition is crucial for mismatch repair. MutS alpha may also play a role in DNA homologous recombination repair. In melanocytes may modulate both UV-B-induced cell cycle regulation and apoptosis.
Cellular Location	Nucleus. Chromosome
Tissue Location	Ubiquitously expressed.

Images



Western blot analysis of MSH2 expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.