

Aspartate Aminotransferase Antibody

Rabbit mAb

Catalog # AP91566

Product Information

Application	WB, IHC, IF, FC, ICC, IHF
Primary Accession	P17174
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	Aspartate aminotransferase 1; ASTQTL1; cAspAT; cCAT; GIG18; Glutamate oxaloacetate transaminase 1; GOT1; Growth inhibiting protein 18; SGOT; Transaminase A;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	46248


Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 FC 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Aspartate Aminotransferase
Description	Plays a key role in amino acid metabolism.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	GOT1 (HGNC:4432)
Function	Biosynthesis of L-glutamate from L-aspartate or L-cysteine (PubMed: 21900944). Important regulator of levels of glutamate, the major excitatory neurotransmitter of the vertebrate central nervous system. Acts as a scavenger of glutamate in brain neuroprotection. The aspartate aminotransferase activity is involved in hepatic glucose synthesis during development and in adipocyte glyceroneogenesis. Using L-cysteine as substrate, regulates levels of mercaptopyruvate, an important source of hydrogen sulfide. Mercaptopyruvate is converted into H(2)S via the action of 3-mercaptopyruvate sulfurtransferase (3MST). Hydrogen sulfide is an important synaptic modulator and neuroprotectant in the brain. In addition, catalyzes (2S)-2- aminobutanoate, a by-product in the cysteine biosynthesis pathway (PubMed: 27827456).
Cellular Location	Cytoplasm.

Images



Western blot analysis of Aspartate Aminotransferase expression in MCF7 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.