

Mad2L2 Antibody

Rabbit mAb

Catalog # AP91576

Product Information

Application	WB, IHC, IF, FC, ICC, IP, IHF
Primary Accession	Q9UI95
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	hREV7; MAD2B; Mad2L2 Mitotic Arrest Deficient 2 L2; POLZ2; REV 7; REV7; REV7 homolog;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	24334

Additional Information

Dilution	WB 1:500~1:1000 IHC 1:100~1:500 ICC/IF 1:50~1:200 IP 1:30 FC 1:40
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Mad2L2
Description	Adapter protein able to interact with different proteins and involved in different biological processes. Mediates the interaction between the error-prone DNA polymerase zeta catalytic subunit REV3L and the inserter polymerase REV1, thereby mediating the second polymerase switching in translesion DNA synthesis.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	MAD2L2
Synonyms	MAD2B, REV7
Function	Adapter protein able to interact with different proteins and involved in different biological processes (PubMed: 11459825 , PubMed: 11459826 , PubMed: 17296730 , PubMed: 17719540 , PubMed: 19443654 , PubMed: 29656893). Mediates the interaction between the error-prone DNA polymerase zeta catalytic subunit REV3L and the inserter polymerase REV1, thereby mediating the second polymerase switching in translesion DNA synthesis (PubMed: 20164194). Translesion DNA synthesis releases the replication blockade of replicative polymerases, stalled in presence of DNA lesions (PubMed: 20164194). Component of the shieldin complex, which plays an important role in repair of DNA double-stranded breaks (DSBs) (PubMed: 29656893). During G1 and S phase of the cell cycle, the complex

functions downstream of TP53BP1 to promote non-homologous end joining (NHEJ) and suppress DNA end resection (PubMed:[29656893](#)). Mediates various NHEJ-dependent processes including immunoglobulin class-switch recombination, and fusion of unprotected telomeres (PubMed:[29656893](#)). May also regulate another aspect of cellular response to DNA damage through regulation of the JNK-mediated phosphorylation and activation of the transcriptional activator ELK1 (PubMed:[17296730](#)). Inhibits the FZR1- and probably CDC20-mediated activation of the anaphase promoting complex APC thereby regulating progression through the cell cycle (PubMed:[11459825](#), PubMed:[17719540](#)). Regulates TCF7L2-mediated gene transcription and may play a role in epithelial-mesenchymal transdifferentiation (PubMed:[19443654](#)).


Cellular Location

Nucleus. Cytoplasm, cytoskeleton, spindle. Cytoplasm. Chromosome.
Note=Recruited to sites of chromosomal double-stranded breaks during G1 and S phase of the cell cycle

Tissue Location

Ubiquitously expressed.

Images



Western blot analysis of Mad2L2 expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.