

# TRPM7 Antibody

Rabbit mAb

Catalog # AP91591

## Product Information

<b>Application</b>	WB, IF, FC, ICC
<b>Primary Accession</b>	<a href="#">Q96QT4</a>
<b>Reactivity</b>	Rat, Human, Mouse
<b>Clonality</b>	Monoclonal
<b>Other Names</b>	ALSPDC; CHAK; CHAK1; LTrpC7; TRPM7;
<b>Isotype</b>	Rabbit IgG
<b>Host</b>	Rabbit
<b>Calculated MW</b>	212697

## Additional Information

<b>Dilution</b>	WB 1:500~1:2000 ICC/IF 1:50~1:200 FC 1:100
<b>Purification</b>	Affinity-chromatography
<b>Immunogen</b>	A synthesized peptide derived from human TRPM7
<b>Description</b>	Essential ion channel and serine/threonine-protein kinase. Divalent cation channel permeable to calcium and magnesium. Has a central role in magnesium ion homeostasis and in the regulation of anoxic neuronal cell death.
<b>Storage Condition and Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

## Protein Information

<b>Name</b>	TRPM7
<b>Synonyms</b>	CHAK1, LTRPC7 {ECO:0000303   PubMed:113855}
<b>Function</b>	Bifunctional protein that combines an ion channel with an intrinsic kinase domain, enabling it to modulate cellular functions either by conducting ions through the pore or by phosphorylating downstream proteins via its kinase domain. The channel is highly permeable to divalent cations, specifically calcium (Ca <sup>2+</sup> ), magnesium (Mg <sup>2+</sup> ) and zinc (Zn <sup>2+</sup> ) and mediates their influx (PubMed: <a href="#">11385574</a> , PubMed: <a href="#">12887921</a> , PubMed: <a href="#">15485879</a> , PubMed: <a href="#">24316671</a> , PubMed: <a href="#">35561741</a> , PubMed: <a href="#">36027648</a> ). Controls a wide range of biological processes such as Ca <sup>2+</sup> (+), Mg <sup>2+</sup> (+) and Zn <sup>2+</sup> (+) homeostasis, vesicular Zn <sup>2+</sup> (+) release channel and intracellular Ca <sup>2+</sup> (+) signaling, embryonic development, immune responses, cell motility, proliferation and differentiation (By similarity). The C-terminal alpha-kinase domain autophosphorylates cytoplasmic residues of TRPM7 (PubMed: <a href="#">18365021</a> ). In vivo, TRPM7 phosphorylates SMAD2, suggesting that TRPM7 kinase may play a


role in activating SMAD signaling pathways. In vitro, TRPM7 kinase phosphorylates ANXA1 (annexin A1), myosin II isoforms and a variety of proteins with diverse cellular functions (PubMed:[15485879](#), PubMed:[18394644](#)).

### Cellular Location

Cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q923J1}. Cytoplasmic vesicle membrane {ECO:0000250|UniProtKB:Q923J1}; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q923J1}. Note=Localized largely in intracellular Zn(2+)-storage vesicles. {ECO:0000250|UniProtKB:Q923J1}

### Images

---



Western blot analysis of TRPM7 expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.