

# DNA Ligase IV Antibody

Rabbit mAb

Catalog # AP91608

## Product Information

<b>Application</b>	WB, IHC, IF, ICC, IHF
<b>Primary Accession</b>	<a href="#">P49917</a>
<b>Reactivity</b>	Human
<b>Clonality</b>	Monoclonal
<b>Other Names</b>	DNA joinase; DNA ligase 4; DNA ligase IV; DNA repair enzyme; LIG4; LIG4S; Ligase IV;
<b>Isotype</b>	Rabbit IgG
<b>Host</b>	Rabbit
<b>Calculated MW</b>	103971

## Additional Information

<b>Dilution</b>	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200
<b>Purification</b>	Affinity-chromatography
<b>Immunogen</b>	A synthesized peptide derived from human DNA Ligase IV
<b>Description</b>	Efficiently joins single-strand breaks in a double-stranded polydeoxynucleotide in an ATP-dependent reaction. Involved in DNA non-homologous end joining (NHEJ) required for double-strand break repair and V(D)J recombination.
<b>Storage Condition and Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

## Protein Information

<b>Name</b>	LIG4 {ECO:0000303   PubMed:16357942, ECO:0000312   HGNC:HGNC:6601}
<b>Function</b>	DNA ligase involved in DNA non-homologous end joining (NHEJ); required for double-strand break (DSB) repair and V(D)J recombination (PubMed: <a href="#">12517771</a> , PubMed: <a href="#">17290226</a> , PubMed: <a href="#">23523427</a> , PubMed: <a href="#">29980672</a> , PubMed: <a href="#">33586762</a> , PubMed: <a href="#">8798671</a> , PubMed: <a href="#">9242410</a> , PubMed: <a href="#">9809069</a> ). Catalyzes the NHEJ ligation step of the broken DNA during DSB repair by resealing the DNA breaks after the gap filling is completed (PubMed: <a href="#">12517771</a> , PubMed: <a href="#">17290226</a> , PubMed: <a href="#">9242410</a> , PubMed: <a href="#">9809069</a> ). Joins single-strand breaks in a double-stranded polydeoxynucleotide in an ATP-dependent reaction (PubMed: <a href="#">12517771</a> , PubMed: <a href="#">17290226</a> , PubMed: <a href="#">9242410</a> , PubMed: <a href="#">9809069</a> ). LIG4 is mechanistically flexible: it can ligate nicks as well as compatible DNA overhangs alone, while in the presence of XRCC4, it can ligate ends with 2-nucleotides (nt) microhomology and 1-nt gaps (PubMed: <a href="#">17290226</a> ). Forms a subcomplex with XRCC4; the LIG4-XRCC4 subcomplex is responsible for the NHEJ ligation step and XRCC4 enhances the

joining activity of LIG4 (PubMed:[9242410](#), PubMed:[9809069](#)). Binding of the LIG4-XRCC4 complex to DNA ends is dependent on the assembly of the DNA-dependent protein kinase complex DNA-PK to these DNA ends (PubMed:[10854421](#)). LIG4 regulates nuclear localization of XRCC4 (PubMed:[24984242](#)).

**Cellular Location**


Nucleus

**Tissue Location**

Testis, thymus, prostate and heart.

## Images

---



Western blot analysis of DNA Ligase IV expression in Ramos cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.