

# TMEM173 Antibody

Rabbit mAb

Catalog # AP91634

## Product Information

---

<b>Application</b>	WB, IF, FC, ICC
<b>Primary Accession</b>	<a href="#">Q86WV6</a>
<b>Reactivity</b>	Human
<b>Clonality</b>	Monoclonal
<b>Other Names</b>	ERIS; hMITA; hSTING; MITA; MPYS; NET23; STING; Tmem173; STING1 ; Stimulator of interferon genes protein ;
<b>Isotype</b>	Rabbit IgG
<b>Host</b>	Rabbit
<b>Calculated MW</b>	42193

## Additional Information

---

<b>Dilution</b>	WB 1:500~1:2000 ICC/IF 1:100~1:500 FC 1:20
<b>Purification</b>	Affinity-chromatography
<b>Immunogen</b>	A synthesized peptide derived from human TMEM173
<b>Description</b>	Facilitator of innate immune signaling that promotes the production of type I interferon (IFN-alpha and IFN-beta). Innate immune response is triggered in response to non-CpG double-stranded DNA from viruses and bacteria delivered to the cytoplasm.
<b>Storage Condition and Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

## Protein Information

---

<b>Name</b>	<a href="#">STING1 ( HGNC:27962)</a>
<b>Function</b>	Facilitator of innate immune signaling that acts as a sensor of cytosolic DNA from bacteria and viruses and promotes the production of type I interferon (IFN-alpha and IFN-beta) (PubMed: <a href="#">18724357</a> , PubMed: <a href="#">18818105</a> , PubMed: <a href="#">19433799</a> , PubMed: <a href="#">19776740</a> , PubMed: <a href="#">23027953</a> , PubMed: <a href="#">23747010</a> , PubMed: <a href="#">23910378</a> , PubMed: <a href="#">27801882</a> , PubMed: <a href="#">29973723</a> , PubMed: <a href="#">30842659</a> , PubMed: <a href="#">35045565</a> , PubMed: <a href="#">35388221</a> , PubMed: <a href="#">36808561</a> , PubMed: <a href="#">37832545</a> , PubMed: <a href="#">25704810</a> , PubMed: <a href="#">39255680</a> ). Innate immune response is triggered in response to non-CpG double-stranded DNA from viruses and bacteria delivered to the cytoplasm (PubMed: <a href="#">26300263</a> ). Acts by binding cyclic dinucleotides: recognizes and binds cyclic di-GMP (c- di-GMP), a second messenger produced by bacteria, cyclic UMP-AMP (2',3'-cUAMP), and cyclic GMP-AMP (cGAMP), a messenger produced by CGAS in response to DNA virus in the cytosol (PubMed: <a href="#">21947006</a> , PubMed: <a href="#">23258412</a> , PubMed: <a href="#">23707065</a> ,

PubMed:[23722158](#), PubMed:[23747010](#), PubMed:[23910378](#),  
PubMed:[26229117](#), PubMed:[30842659](#), PubMed:[35388221](#),  
PubMed:[37379839](#)). Upon binding to c-di-GMP, cUAMP or cGAMP, STING1 oligomerizes, translocates from the endoplasmic reticulum and is phosphorylated by TBK1 on the pLXIS motif, leading to recruitment and subsequent activation of the transcription factor IRF3 to induce expression of type I interferon and exert a potent anti-viral state (PubMed:[22394562](#),  
PubMed:[25636800](#), PubMed:[29973723](#), PubMed:[30842653](#)  
PubMed:[35045565](#), PubMed:[35388221](#)). Exhibits 2',3' phosphodiester linkage-specific ligand recognition: can bind both 2'-3' linked cGAMP (2'-3'-cGAMP) and 3'-3' linked cGAMP but is preferentially activated by 2'-3' linked cGAMP (PubMed:[23747010](#), PubMed:[23910378](#), PubMed:[26300263](#)). The preference for 2'-3'-cGAMP, compared to other linkage isomers is probably due to the ligand itself, whichs adopts an organized free- ligand conformation that resembles the STING1-bound conformation and pays low energy costs in changing into the active conformation (PubMed:[26150511](#)). In addition to promote the production of type I interferons, plays a direct role in autophagy (PubMed:[30568238](#), PubMed:[30842662](#)). Following cGAMP-binding, STING1 buds from the endoplasmic reticulum into COPII vesicles, which then form the endoplasmic reticulum-Golgi intermediate compartment (ERGIC) (PubMed:[30842662](#)). The ERGIC serves as the membrane source for WIPI2 recruitment and LC3 lipidation, leading to formation of autophagosomes that target cytosolic DNA or DNA viruses for degradation by the lysosome (PubMed:[30842662](#)). Promotes autophagy by acting as a proton channel that directs proton efflux from the Golgi to facilitate MAP1LC3B/LC3B lipidation (PubMed:[37535724](#)). The autophagy- and interferon-inducing activities can be uncoupled and autophagy induction is independent of TBK1 phosphorylation (PubMed:[30568238](#),  
PubMed:[30842662](#)). Autophagy is also triggered upon infection by bacteria: following c-di-GMP-binding, which is produced by live Gram- positive bacteria, promotes reticulophagy (By similarity). May be involved in translocon function, the translocon possibly being able to influence the induction of type I interferons (PubMed:[18724357](#)). May be involved in transduction of apoptotic signals via its association with the major histocompatibility complex class II (MHC-II) (By similarity). Involved in intercellular immune signaling. Cross-activated by 2',3'- cGAMP previously generated in virus-infected cells, triggers type I interferon signaling in macrophages and uninfected neighboring cells to propagate and amplify the antiviral immune response.

## Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein {ECO:0000255, ECO:0000269 | PubMed:[30842659](#),  
ECO:0000269 | PubMed:[32690950](#)}. Cytoplasm, perinuclear region. Endoplasmic reticulum-Golgi intermediate compartment membrane; Multi-pass membrane protein {ECO:0000255, ECO:0000269 | PubMed:[32690950](#)}. Golgi apparatus membrane; Multi-pass membrane protein. Cytoplasmic vesicle, autophagosome membrane; Multi-pass membrane protein. Mitochondrion outer membrane; Multi-pass membrane protein. Cell membrane {ECO:0000250 | UniProtKB:Q3TBT3}; Multi-pass membrane protein. Note=In response to double-stranded DNA stimulation, translocates from the endoplasmic reticulum through the endoplasmic reticulum-Golgi intermediate compartment and Golgi to post-Golgi vesicles, where the kinase TBK1 is recruited (PubMed:[19433799](#), PubMed:[29694889](#), PubMed:[30842653](#), PubMed:[30842659](#)). Upon cGAMP-binding, translocates to the endoplasmic reticulum-Golgi intermediate compartment (ERGIC) in a process that is dependent on COPII vesicles; STING1-containing ERGIC serves as a membrane source for LC3 lipidation, which is a key step in autophagosome biogenesis (PubMed:[30842662](#), PubMed:[37832545](#)). Localizes in the lysosome membrane in a TMEM203-dependent manner (By similarity). {ECO:0000250 | UniProtKB:Q3TBT3, ECO:0000269 | PubMed:[19433799](#), ECO:0000269 | PubMed:[29694889](#),


ECO:0000269 | PubMed:30842653, ECO:0000269 | PubMed:30842659,  
ECO:0000269 | PubMed:30842662, ECO:0000269 | PubMed:32690950,  
ECO:0000269 | PubMed:37832545}

#### Tissue Location

Ubiquitously expressed (PubMed:18724357, PubMed:18818105). Expressed in skin endothelial cells, alveolar type 2 pneumocytes, bronchial epithelium and alveolar macrophages (PubMed:25029335).

#### Images

---



Western blot analysis of TMEM173 expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.