

# DEK Antibody

Rabbit mAb

Catalog # AP91703

## Product Information

---

|                          |                                 |
|--------------------------|---------------------------------|
| <b>Application</b>       | WB, IHC, IF, ICC, IP, IHF       |
| <b>Primary Accession</b> | <a href="#">P35659</a>          |
| <b>Reactivity</b>        | Rat, Human                      |
| <b>Clonality</b>         | Monoclonal                      |
| <b>Other Names</b>       | Dek; DEK oncogene; Protein DEK; |
| <b>Isotype</b>           | Rabbit IgG                      |
| <b>Host</b>              | Rabbit                          |
| <b>Calculated MW</b>     | 42674                           |

## Additional Information

---

|                                     |   |
|-------------------------------------|---|
| <b>Dilution</b>                     | WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 IP 1:50  |
| <b>Purification</b>                 | Affinity-chromatography   |
| <b>Immunogen</b>                    | A synthesized peptide derived from human DEK  |
| <b>Description</b>                  | Involved in chromatin organization.   |
| <b>Storage Condition and Buffer</b> | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |

## Protein Information


---

|                          |   |
|--------------------------|---|
| <b>Name</b>              | DEK   |
| <b>Function</b>          | Involved in chromatin organization.   |
| <b>Cellular Location</b> | Nucleus. Note=Enriched in regions where chromatin is decondensed or sparse in the interphase nuclei |
| <b>Tissue Location</b>   | Ubiquitous. Expressed at relatively high levels.  |

## Images

---

Western blot analysis of DEK expression in 293T cell lysate.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.