

TRP1 Antibody

Rabbit mAb

Catalog # AP91747

Product Information

Application	WB, IHC, IP
Primary Accession	P17643
Reactivity	Human
Clonality	Monoclonal
Other Names	CAS2; CATB; GP75; OCA3; TRP1; TYRP; TYRP1; TYRRP;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	60724


Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200 IP 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human TRP1
Description	Oxidation of 5,6-dihydroxyindole-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid. May regulate or influence the type of melanin synthesized.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	TYRP1 (HGNC:12450)
Function	Plays a role in melanin biosynthesis (PubMed: 16704458 , PubMed: 22556244 , PubMed: 23504663). Catalyzes the oxidation of 5,6-dihydroxyindole-2-carboxylic acid (DHICA) into indole-5,6-quinone-2-carboxylic acid in the presence of bound Cu(2+) ions, but not in the presence of Zn(2+) (PubMed: 28661582). May regulate or influence the type of melanin synthesized (PubMed: 16704458 , PubMed: 22556244). Also to a lower extent, capable of hydroxylating tyrosine and producing melanin (By similarity).
Cellular Location	Melanosome membrane {ECO:0000250 UniProtKB:P07147}; Single-pass type I membrane protein {ECO:0000250 UniProtKB:P07147}. Note=Located to mature stage III and IV melanosomes and apposed endosomal tubular membranes. Transported to pigmented melanosomes by the BLOC-1 complex. Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38 {ECO:0000250 UniProtKB:P07147}
Tissue Location	Pigment cells.

Images



Western blot analysis of TRP1 expression in Human melanoma lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.