

VPS26 Antibody

Rabbit mAb

Catalog # AP91752

Product Information

Application	WB, IF, ICC, IP
Primary Accession	O75436
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	HB58; Hbeta58; hVPS26; PEP8A; VPS 26; VPS26A;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	38170

Additional Information

Dilution	WB 1:500~1:2000 ICC/IF 1:50~1:200 IP 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human VPS26
Description	Essential component of the retromer complex, a complex required to retrieve lysosomal enzyme receptors (IGF2R and M6PR) from endosomes to the trans-Golgi network. Also required to regulate transcytosis of the polymeric immunoglobulin receptor (pIgR-pIgA).
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information


Name	VPS26A {ECO:0000303 PubMed:30213940, ECO:0000312 HGNC:HGNC:12711}
Function	Acts as a component of the retromer cargo-selective complex (CSC). The CSC is believed to be the core functional component of retromer or respective retromer complex variants acting to prevent missorting of selected transmembrane cargo proteins into the lysosomal degradation pathway. The recruitment of the CSC to the endosomal membrane involves RAB7A and SNX3. The SNX-BAR retromer mediates retrograde transport of cargo proteins from endosomes to the trans- Golgi network (TGN) and is involved in endosome-to-plasma membrane transport for cargo protein recycling. The SNX3-retromer mediates the retrograde endosome-to-TGN transport of WLS distinct from the SNX-BAR retromer pathway. The SNX27-retromer is believed to be involved in endosome-to-plasma membrane trafficking and recycling of a broad spectrum of cargo proteins (Probable). The CSC seems to act as recruitment hub for other proteins, such as the WASH complex and TBC1D5 (Probable). Required for retrograde transport of lysosomal enzyme receptor

IGF2R (PubMed:[15078902](#), PubMed:[15078903](#)). Required to regulate transcytosis of the polymeric immunoglobulin receptor (pIgR-pIgA) (PubMed:[15247922](#)). Required for the endosomal localization of WASHC2A (indicative for the WASH complex) (PubMed:[22070227](#)). Required for the endosomal localization of TBC1D5 (PubMed:[20923837](#)). Mediates retromer cargo recognition of SORL1 and is involved in trafficking of SORL1 implicated in sorting and processing of APP (PubMed:[22279231](#)). Involved in retromer-independent lysosomal sorting of F2R (PubMed:[16407403](#)). Involved in recycling of ADRB2 (PubMed:[21602791](#)). Enhances the affinity of SNX27 for PDZ-binding motifs in cargo proteins (By similarity).

Cellular Location

Cytoplasm. Endosome membrane; Peripheral membrane protein {ECO:0000250|UniProtKB:P40336}. Early endosome Note=Localizes to tubular profiles adjacent to endosomes (PubMed:15078903). Predominantly found in early not late endosomes (By similarity). {ECO:0000250|UniProtKB:P40336}

Images



Western blot analysis of VPS26 expression in A431 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.