

CoREST Antibody

Rabbit mAb

Catalog # AP91767

Product Information

Application	WB, IHC, IF, ICC, IHF
Primary Accession	Q9UKL0
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	COREST; Nr2b2; RCOR; Rcor1; Rxrb;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	53327

Additional Information

Dilution	WB 1:1000~1:5000 IHC 1:50~1:200 ICC/IF 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human CoREST
Description	Essential component of the BHC complex, a corepressor complex that represses transcription of neuron-specific genes in non-neuronal cells. The BHC complex is recruited at RE1/NRSE sites by REST and acts by deacetylating and demethylating specific sites on histones, thereby acting as a chromatin modifier.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	RCOR1
Synonyms	KIAA0071, RCOR
Function	Essential component of the BHC complex, a corepressor complex that represses transcription of neuron-specific genes in non-neuronal cells. The BHC complex is recruited at RE1/NRSE sites by REST and acts by deacetylating and demethylating specific sites on histones, thereby acting as a chromatin modifier. In the BHC complex, it serves as a molecular beacon for the recruitment of molecular machinery, including MeCP2 and SUV39H1, that imposes silencing across a chromosomal interval. Plays a central role in demethylation of Lys-4 of histone H3 by promoting demethylase activity of KDM1A on core histones and nucleosomal substrates. It also protects KDM1A from the proteasome. Component of a RCOR/GFI/KDM1A/HDAC complex that suppresses, via histone deacetylase (HDAC) recruitment, a number of genes implicated in multilineage blood cell development and controls hematopoietic

differentiation.


Cellular Location

Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00512, ECO:0000255 | PROSITE-ProRule:PRU00624, ECO:0000269 | PubMed:10734093, ECO:0000269 | PubMed:15897453}. Note=Upon infection by HSV-1, it is partially translocated into the cytoplasm in an HSV-1-dependent manner

Tissue Location

Ubiquitously expressed.

Images



Western blot analysis of CoREST expression in Molt-4 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.