

NEDD4 Antibody

Rabbit mAb

Catalog # AP91782

Product Information

Application	WB, FC, IP
Primary Accession	P46934
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	Nedd4; PIG53; RPF1;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	149114

Additional Information

Dilution	WB 1:1000~1:5000 IP 1:50 FC 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human NEDD4
Description	E3 ubiquitin-protein ligase which accepts ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates. Involved in the pathway leading to the degradation of VEGFR-2/KDFR, independently of its ubiquitin-ligase activity.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information


Name	NEDD4
Synonyms	KIAA0093, NEDD4-1, RPF1 {ECO:0000303 Pub
Function	E3 ubiquitin-protein ligase which accepts ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates. Specifically ubiquitinates 'Lys-63' in target proteins (PubMed: 19920177 , PubMed: 21399620 , PubMed: 23644597). Involved in the pathway leading to the degradation of VEGFR-2/KDFR, independently of its ubiquitin-ligase activity. Monoubiquitinates IGF1R at multiple sites, thus leading to receptor internalization and degradation in lysosomes (By similarity). Ubiquitinates FGFR1, leading to receptor internalization and degradation in lysosomes (PubMed: 21765395). Promotes ubiquitination of RAPGEF2 (PubMed: 11598133). According to PubMed: 18562292 the direct link between NEDD4 and PTEN regulation through polyubiquitination described in

PubMed:[17218260](#) is questionable. Involved in ubiquitination of ERBB4 intracellular domain E4ICD (By similarity). Part of a signaling complex composed of NEDD4, RAP2A and TNIK which regulates neuronal dendrite extension and arborization during development (By similarity). Ubiquitinates TNK2 and regulates EGF-induced degradation of EGFR and TNF2 (PubMed:[20086093](#)). Ubiquitinates BRAT1 and this ubiquitination is enhanced in the presence of NDFIP1 (PubMed:[25631046](#)). Ubiquitinates DAZAP2, leading to its proteasomal degradation (PubMed:[11342538](#)). Ubiquitinates POLR2A (PubMed:[19920177](#)). Functions as a platform to recruit USP13 to form an NEDD4-USP13 deubiquitination complex that plays a critical role in cleaving the 'Lys-48'-linked ubiquitin chains of VPS34 and then stabilizing VPS34, thus promoting the formation of autophagosomes (PubMed:[32101753](#)).

Cellular Location

Cytoplasm. Nucleus. Cell membrane {ECO:0000250 | UniProtKB:P46935}; Peripheral membrane protein {ECO:0000250 | UniProtKB:P46935}. Note=Predominantly cytoplasmic but also located in the nucleus (PubMed:11342538). Recruited to the plasma membrane by GRB10. Once complexed with GRB10 and IGF1R, follows IGF1R internalization, remaining associated with early endosomes. Uncouples from IGF1R-containing endosomes before the sorting of the receptor to the lysosomal compartment (By similarity). May be recruited to exosomes by NDFIP1 (PubMed:18819914). {ECO:0000250 | UniProtKB:P46935, ECO:0000269 | PubMed:11342538, ECO:0000269 | PubMed:18819914}

Images



Western blot analysis of NEDD4 expression in A549 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.