

# **SAE1 Antibody**

Rabbit mAb Catalog # AP91797

#### **Product Information**

**Application** WB, IF, FC, ICC, IP

Primary Accession

Reactivity

Clonality

Q9UBE0

Rat, Human

Monoclonal

Other Names AOS1; HSPC140; Sae1; SUA1; UBLE1A;

IsotypeRabbit IgGHostRabbitCalculated MW38450

#### **Additional Information**

**Dilution** WB 1:1000~1:5000 ICC/IF 1:50~1:200 IP 1:50 FC 1:100

**Purification** Affinity-chromatography

**Immunogen** A synthesized peptide derived from human SAE1

**Description** The heterodimer acts as a E1 ligase for SUMO1, SUMO2, SUMO3, and

probably SUMO4. It mediates ATP-dependent activation of SUMO proteins followed by formation of a thioester bond between a SUMO protein and a

conserved active site cysteine residue on UBA2/SAE2.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

## **Protein Information**

Name SAE1

**Synonyms** AOS1, SUA1, UBLE1A

**Function** The heterodimer acts as an E1 ligase for SUMO1, SUMO2, SUMO3, and

probably SUMO4. It mediates ATP-dependent activation of SUMO proteins followed by formation of a thioester bond between a SUMO protein and a


conserved active site cysteine residue on UBA2/SAE2.

Cellular Location Nucleus.

**Tissue Location** Expression level increases during S phase and drops in G2 phase (at protein

level).

### **Images**



Western blot analysis of SAE1 expression in Jurkat cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.