10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

SOCS2 Antibody

Rabbit mAb
Catalog # AP91836

Product Information

Application	WB, IF, ICC, IP
Primary Accession	O14508
Reactivity	Human
Clonality	Monoclonal
Other Names	CIS2; Cish2; Socs2; SSI2; STATI2;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	22172

Additional Information

Dilution	WB 1:500~1:2000 ICC/IF 1:50~1:200 IP 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human SOCS2
Description	SOCS family proteins form part of a classical negative feedback system that regulates cytokine signal transduction. SOCS2 appears to be a negative regulator in the growth hormone/IGF1 signaling pathway.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	SOCS2 {ECO:0000303|PubMed:10512686, ECO:0000312|HGNC:HGNC:19382}
Function	Substrate-recognition component of a cullin-5-RING E3 ubiquitin-protein ligase complex (ECS complex, also named CRL5 complex), which mediates the ubiquitination and subsequent proteasomal degradation of target proteins, such as EPOR and GHR (PubMed:11781573, PubMed:21980433, PubMed:25505247, PubMed:31182716, PubMed:34857742). Specifically recognizes and binds phosphorylated proteins via its SH2 domain, promoting their ubiquitination (PubMed:21980433, PubMed:25505247, PubMed:31182716, PubMed:34857742, PubMed:37816714). The ECS(SOCS2) complex acts as a key regulator of growth hormone receptor (GHR) levels by mediating ubiquitination and degradation of GHR, following GHR phosphorylation by JAK2 (PubMed:21980433, PubMed:25505247, PubMed:34857742). The ECS(SOCS2) also catalyzes ubiquitination and degradation of JAK2-phosphorylated EPOR (PubMed:11781573).
Cellular Location	Cytoplasm.

Tissue Location	High expression in heart, placenta, lung, kidney and prostate. Predominantly expressed in pulmonary epithelia cells, specifically type II pneumocytes.

Images



Western blot analysis of SOCS2 expression in K562 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.