10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

NM23A Antibody

Rabbit mAb
Catalog # AP91847

Product Information

Application	WB, IHC, IF, FC, ICC, IHF
Primary Accession	P15531
Reactivity	Human
Clonality	Monoclonal
Other Names	NB; AWD; NBS; GAAD; NM23; NDPKA; NDPK-A; NM23-H1; NME1;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	17149

Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 FC 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human NM23A
Description	Major role in the synthesis of nucleoside triphosphates other than ATP. Possesses nucleoside-diphosphate kinase, serine/threonine-specific protein kinase, geranyl and farnesyl pyrophosphate kinase, histidine protein kinase and 3'-5' exonuclease activities.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	NME1 (HGNC:7849)
Function	Catalyzes the transfer of a gamma-phosphoryl group from a nucleoside triphosphate, mainly ATP, to a nucleoside diphosphate via a ping-pong mechanism involving a phosphohistidine intermediate, therefore contributing to the nucleoside triphosphate homeostasis (PubMed:10952986, PubMed:14960567, PubMed:16313181, PubMed:1851158, PubMed:23519676, PubMed:33903070, PubMed:8810265, PubMed:9038158). Also phosphorylates geranyl pyrophosphate (GPP) and farnesyl pyrophosphate (FPP), linking it to isoprenoid metabolism (PubMed:10952986). Additionally, functions as a non-specific serine/threonine kinase and histidine protein kinase, transferring phosphoryl groups from its active site to target proteins (PubMed:8529641, PubMed:9038158). May function as a Mg(2+)-dependent single-stranded DNA endonuclease as part of the SET complex, cooperating with the 3'-5' exonuclease TREX1 to mediate apoptotic DNA fragmentation in cytotoxic T lymphocytes (PubMed:12628186, PubMed:16818237). Reported to nick one DNA strand, enabling TREX1 to remove nucleotides from the free 3' end,

enhancing DNA damage and suppressing DNA end reannealing and repair (PubMed:16818237). Has been shown to cleave double strands DNA within the 3'-portions of both 5'-SHS silencer and NHE basal promoter element of the PDGFA gene, potentially repressing its transcription (PubMed:11694515). May also function as a Mg(2+)-dependent 3'-5' DNA exonuclease, excising nucleotides from 3' single-stranded DNA or DNA with 3' single strand overhangs, suggesting a role in DNA nucleolytic processing (PubMed:14960567, PubMed:16313181). Involved in the regulation of tumor metastasis and cellular differentiation (By similarity). Also required for cell motility (PubMed:8270257, PubMed:25582197). May control, with NME2, AcCoA usage between histone acetylation and fatty acid synthesis, possibly by binding and releasing AcCoA at transcriptionally active chromatin regions in proximity to histone acetyltransferase (HAT) (By similarity).


Cellular Location

Cytoplasm. Nucleus. Cell membrane {ECO:0000250|UniProtKB:P52175}. Note=Cell-cycle dependent nuclear localization which can be induced by degradation of the SET complex by GzmA (PubMed:12628186). In response to DNA damage, translocates to the nucleus where it might participate in DNA nucleolytic processing (PubMed:16313181).

Tissue Location

Ubiquitously expressed (PubMed:12601555, PubMed:16442775). Expressed in tumor cell lines (PubMed:10512675, PubMed:16442775).

Images



Western blot analysis of NM23A expression in Jurkat cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.