

# IGFBP2 Antibody

Rabbit mAb

Catalog # AP91860

## Product Information

<b>Application</b>	WB, IHC
<b>Primary Accession</b>	<a href="#">P18065</a>
<b>Reactivity</b>	Human
<b>Clonality</b>	Monoclonal
<b>Other Names</b>	BP2; IBP2; IGF BP53; IGFBP2; IGFBP53;
<b>Isotype</b>	Rabbit IgG
<b>Host</b>	Rabbit
<b>Calculated MW</b>	34814

## Additional Information

<b>Dilution</b>	WB 1:1000~1:5000 IHC 1:50~1:200
<b>Purification</b>	Affinity-chromatography
<b>Immunogen</b>	A synthesized peptide derived from human IGFBP2
<b>Description</b>	Inhibits IGF-mediated growth and developmental rates. IGF-binding proteins prolong the half-life of the IGFs and have been shown to either inhibit or stimulate the growth promoting effects of the IGFs on cell culture. They alter the interaction of IGFs with their cell surface receptors.
<b>Storage Condition and Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

## Protein Information

<b>Name</b>	IGFBP2
<b>Synonyms</b>	BP2, IBP2
<b>Function</b>	Multifunctional protein that plays a critical role in regulating the availability of IGFs such as IGF1 and IGF2 to their receptors and thereby regulates IGF-mediated cellular processes including proliferation, differentiation, and apoptosis in a cell-type specific manner (PubMed: <a href="#">18563800</a> , PubMed: <a href="#">38796567</a> ). Functions coordinately with receptor protein tyrosine phosphatase beta/PTPRB and the IGF1 receptor to regulate IGF1-mediated signaling by stimulating the phosphorylation of PTEN leading to its inactivation and AKT1 activation (PubMed: <a href="#">22869525</a> ). Plays a positive role in cell migration via interaction with integrin alpha5/ITGA5 through an RGD motif (PubMed: <a href="#">16569642</a> ). Additionally, interaction with ITGA5/ITGB1 enhances the adhesion of endothelial progenitor cells to endothelial cells (PubMed: <a href="#">26076738</a> ). Upon mitochondrial damage, facilitates apoptosis with ITGA5 of podocytes, and then activates the phosphorylation of focal adhesion


kinase (FAK)-mediated mitochondrial injury (PubMed:[38796567](#)).

## Cellular Location

Secreted

## Images

---



Western blot analysis of IGFBP2 expression in SH-SY5Y cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.