

TRF2 Antibody

Rabbit mAb

Catalog # AP91864

Product Information

Application	WB, IHC, IF, ICC, IHF
Primary Accession	Q15554
Reactivity	Human
Clonality	Monoclonal
Other Names	Terf2; TRBF2; TRF2;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	59594

Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human TRF2
Description	Binds the telomeric double-stranded 5'-TTAGGG-3' repeat and plays a central role in telomere maintenance and protection against end-to-end fusion of chromosomes. In addition to its telomeric DNA-binding role, required to recruit a number of factors and enzymes required for telomere protection, including the shelterin complex, TERF2IP/RAP1 and DCLRE1B/Apollo.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	TERF2
Synonyms	TRBF2, TRF2 {ECO:0000303 PubMed:28216226
Function	Binds the telomeric double-stranded 5'-TTAGGG-3' repeat and plays a central role in telomere maintenance and protection against end-to-end fusion of chromosomes (PubMed: 15608617 , PubMed: 16166375 , PubMed: 20655466 , PubMed: 28216226 , PubMed: 9326950 , PubMed: 9326951 , PubMed: 9476899). In addition to its telomeric DNA-binding role, required to recruit a number of factors and enzymes required for telomere protection, including the shelterin complex, TERF2IP/RAP1 and DCLRE1B/Apollo (PubMed: 16166375 , PubMed: 20655466). Component of the shelterin complex (telosome) that is involved in the regulation of telomere length and protection (PubMed: 16166375). Shelterin associates with arrays of double-stranded 5'-TTAGGG-3' repeats added by telomerase and protects chromosome ends; without its protective activity, telomeres are no longer hidden from the DNA

damage surveillance and chromosome ends are inappropriately processed by DNA repair pathways (PubMed:[16166375](#)). Together with DCLRE1B/Apollo, plays a key role in telomeric loop (T loop) formation by generating 3' single-stranded overhang at the leading end telomeres: T loops have been proposed to protect chromosome ends from degradation and repair (PubMed:[20655466](#)). Required both to recruit DCLRE1B/Apollo to telomeres and activate the exonuclease activity of DCLRE1B/Apollo (PubMed:[20655466](#), PubMed:[28216226](#)). Preferentially binds to positive supercoiled DNA (PubMed:[15608617](#), PubMed:[20655466](#)). Together with DCLRE1B/Apollo, required to control the amount of DNA topoisomerase (TOP1, TOP2A and TOP2B) needed for telomere replication during fork passage and prevent aberrant telomere topology (PubMed:[20655466](#)). Recruits TERF2IP/RAP1 to telomeres, thereby participating in to repressing homology-directed repair (HDR), which can affect telomere length (By similarity).


Cellular Location

Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00625, ECO:0000269 | PubMed:20655466}. Chromosome, telomere. Note=Colocalizes with telomeric DNA in interphase cells and is located at chromosome ends during metaphase

Tissue Location

Ubiquitous. Highly expressed in spleen, thymus, prostate, uterus, testis, small intestine, colon and peripheral blood leukocytes.

Images



Western blot analysis of TRF2 expression in Jurkat cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.