

CD98 Antibody

Rabbit mAb

Catalog # AP91865

Product Information

Application	WB, IHC, IP
Primary Accession	P08195
Reactivity	Human
Clonality	Monoclonal
Other Names	4T2HC; CD98; CD98HC; MDU1; NACAE; Slc3a2;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	57945

Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200 IP 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human CD98
Description	Required for the function of light chain amino-acid transporters. Involved in sodium-independent, high-affinity transport of large neutral amino acids such as phenylalanine, tyrosine, leucine, arginine and tryptophan. Involved in guiding and targeting of LAT1 and LAT2 to the plasma membrane.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	SLC3A2 (HGNC:11026)
Synonyms	MDU1
Function	Acts as a chaperone that facilitates biogenesis and trafficking of functional transporters heterodimers to the plasma membrane. Forms heterodimer with SLC7 family transporters (SLC7A5, SLC7A6, SLC7A7, SLC7A8, SLC7A10 and SLC7A11), a group of amino-acid antiporters (PubMed: 10574970 , PubMed: 10903140 , PubMed: 11557028 , PubMed: 30867591 , PubMed: 33298890 , PubMed: 33758168 , PubMed: 34880232 , PubMed: 9751058 , PubMed: 9829974 , PubMed: 9878049). Heterodimers function as amino acids exchangers, the specificity of the substrate depending on the SLC7A subunit. Heterodimers SLC3A2/SLC7A6 or SLC3A2/SLC7A7 mediate the uptake of dibasic amino acids (PubMed: 10903140 , PubMed: 9829974). Heterodimer SLC3A2/SLC7A11 functions as an antiporter by mediating the exchange of extracellular anionic L-cystine and intracellular L-glutamate across the cellular plasma membrane (PubMed: 34880232). SLC3A2/SLC7A10 translocates small

neutral L- and D- amino acids across the plasma membrane (By similarity). SLC3A2/SLC75 or SLC3A2/SLC7A8 translocates neutral amino acids with broad specificity, thyroid hormones and L-DOPA (PubMed:[10574970](#), PubMed:[11389679](#), PubMed:[11557028](#), PubMed:[11564694](#), PubMed:[11742812](#), PubMed:[12117417](#), PubMed:[12225859](#), PubMed:[12716892](#), PubMed:[15980244](#), PubMed:[30867591](#), PubMed:[33298890](#), PubMed:[33758168](#)). SLC3A2 is essential for plasma membrane localization, stability, and the transport activity of SLC7A5 and SLC7A8 (PubMed:[10391915](#), PubMed:[10574970](#), PubMed:[11311135](#), PubMed:[15769744](#), PubMed:[33066406](#)). When associated with LAPT4B, the heterodimer SLC7A5 is recruited to lysosomes to promote leucine uptake into these organelles, and thereby mediates mTORC1 activation (PubMed:[25998567](#)). Modulates integrin-related signaling and is essential for integrin-dependent cell spreading, migration and tumor progression (PubMed:[11121428](#), PubMed:[15625115](#)).


Cellular Location

Apical cell membrane. Cell membrane; Single-pass type II membrane protein. Cell junction {ECO:0000250 | UniProtKB:P10852}. Lysosome membrane. Melanosome. Basolateral cell membrane {ECO:0000250 | UniProtKB:P10852}. Note=Localized at the plasma membrane when associated with SLC7A5/LAT1 or SLC7A8/LAT2 (PubMed:11311135, PubMed:9751058). Localized to the apical membrane of placental syncytiotrophoblastic cells (PubMed:11742812). Recruited to lysosomes by LAPT4B (PubMed:25998567). Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065) Located selectively at cell-cell adhesion sites (By similarity) Colocalized with SLC7A8/LAT2 at the basolateral membrane of kidney proximal tubules and small intestine epithelia. Expressed in both luminal and abluminal membranes of brain capillary endothelial cells (By similarity). {ECO:0000250 | UniProtKB:P10852, ECO:0000269 | PubMed:11311135, ECO:0000269 | PubMed:11742812, ECO:0000269 | PubMed:17081065, ECO:0000269 | PubMed:25998567, ECO:0000269 | PubMed:9751058}

Tissue Location

Expressed ubiquitously in all tissues tested with highest levels detected in kidney, placenta and testis and weakest level in thymus. During gestation, expression in the placenta was significantly stronger at full-term than at the mid-trimester stage. Expressed in HUVECS and at low levels in resting peripheral blood T- lymphocytes and quiescent fibroblasts. Also expressed in fetal liver and in the astrocytic process of primary astrocytic gliomas. Expressed in retinal endothelial cells and in the intestinal epithelial cell line C2BBe1.

Images



Western blot analysis of CD98 expression in HepG2 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.