

Frizzled 9 / CD349 Antibody

Rabbit mAb

Catalog # AP91878

Product Information

| | |
|--------------------------|-------------------------------------|
| Application | WB |
| Primary Accession | O00144 |
| Reactivity | Rat, Human, Mouse |
| Clonality | Monoclonal |
| Other Names | CD349; frizzled-9; Fz-9; FZD3; FzE6 |
| Isotype | Rabbit IgG |
| Host | Rabbit |
| Calculated MW | 64466 |

Additional Information

| | |
|-------------------------------------|---|
| Dilution | WB 1:500~1:2000 |
| Purification | Affinity-chromatography |
| Immunogen | A synthesized peptide derived from human Frizzled 9 / CD349 |
| Description | Receptor for Wnt proteins. Most of frizzled receptors are coupled to the beta-catenin canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes. |
| Storage Condition and Buffer | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |

Protein Information

| | |
|-----------------|---|
| Name | FZD9 |
| Synonyms | FZD3 |
| Function | Receptor for WNT2 that is coupled to the beta-catenin canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes (By similarity). Plays a role in neuromuscular junction (NMJ) assembly by negatively regulating the clustering of acetylcholine receptors (AChR) through the beta-catenin canonical signaling pathway (By similarity). May play a role in neural progenitor cells (NPCs) viability through the beta-catenin canonical signaling pathway by negatively regulating cell cycle arrest leading to inhibition of neuron apoptotic process (PubMed: 27509850). During hippocampal development, regulates neuroblast proliferation and apoptotic cell death. Controls bone formation through non canonical Wnt signaling mediated via ISG15. Positively regulates bone regeneration through non canonical Wnt signaling (By similarity). |


Cellular Location

Cell membrane {ECO:0000250|UniProtKB:Q9R216}; Multi-pass membrane protein. Note=Relocalizes DVL1 to the cell membrane leading to phosphorylation of DVL1 and AXIN1 relocalization to the cell membrane. {ECO:0000250|UniProtKB:Q8K4C8}

Tissue Location

Expressed predominantly in adult and fetal brain, testis, eye, skeletal muscle and kidney. Moderately expressed in pancreas, thyroid, adrenal cortex, small intestine and stomach Detected in fetal liver and kidney. Expressed in neural progenitor cells (PubMed:27509850).

Images



Western blot analysis of Frizzled 9 expression in Human Seminoma lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.