

Prostatic Acid Phosphatase Antibody

Rabbit mAb Catalog # AP91880

Product Information

Application	WB, IHC, IF, ICC, IP, IHF
Primary Accession	<u>P15309</u>
Reactivity	Human, Mouse
Clonality	Monoclonal
Other Names	ACP3; PAP; TMPase;
lsotype	Rabbit IgG
Host	Rabbit
Calculated MW	44566

Additional Information


Dilution Purification Immunogen	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 IP 1:50 Affinity-chromatography A synthesized peptide derived from human Prostatic Acid Phosphatase
Description	A non-specific tyrosine phosphatase that dephosphorylates a diverse number of substrates under acidic conditions (pH 4-6) including alkyl, aryl, and acyl orthophosphate monoesters and phosphorylated proteins. Has lipid phosphatase activity and inactivates lysophosphatidic acid in seminal plasma.
Storage Condition and Buffer	

Protein Information

Name	ACP3 (<u>HGNC:125</u>)
Synonyms	ACPP
Function	A non-specific tyrosine phosphatase that dephosphorylates a diverse number of substrates under acidic conditions (pH 4-6) including alkyl, aryl, and acyl orthophosphate monoesters and phosphorylated proteins (PubMed: <u>10506173</u> , PubMed: <u>15280042</u> , PubMed: <u>20498373</u> , PubMed: <u>9584846</u>). Has lipid phosphatase activity and inactivates lysophosphatidic acid in seminal plasma (PubMed: <u>10506173</u> , PubMed: <u>15280042</u>).
Cellular Location	[Isoform 1]: Secreted
Tissue Location	Highly expressed in the prostate, restricted to glandular and ductal epithelial cells. Also expressed in bladder, kidney, pancreas, lung, cervix, testis and ovary. Weak expression in a subset of pancreatic islet cells, squamous

epithelia, the pilosebaceous unit, colonic neuroendocrine cells and skin adnexal structures. Low expression in prostate carcinoma cells and tissues

Images



Western blot analysis of Prostatic Acid Phosphatase expression in human prostate cancer lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.