

# CD48 Antibody

Rabbit mAb Catalog # AP91882

### **Product Information**

Application	WB, IHC, IF, ICC, IP, IHF
Primary Accession	<u>P09326</u>
Reactivity	Human
Clonality	Monoclonal
Other Names	BCM1; BLAST; BLAST1; CD48; hCD48; mCD48; MEM102; SLAMF2;
lsotype	Rabbit IgG
Host	Rabbit
Calculated MW	27683


#### **Additional Information**

Dilution Purification Immunogen Description	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 IP 1:50 Affinity-chromatography A synthesized peptide derived from human CD48 Ligand for CD2. Might facilitate interaction between activated lymphocytes. Probably involved in regulating T-cell activation.
Storage Condition and Buffer	

#### **Protein Information**

Name	CD48
Synonyms	BCM1, BLAST1
Function	Glycosylphosphatidylinositol (GPI)-anchored cell surface glycoprotein that interacts via its N-terminal immunoglobulin domain with cell surface receptors including CD244/2B4 or CD2 to regulate immune cell function and activation (PubMed:12007789, PubMed:19494291, PubMed:27249817, PubMed:9841922). Participates in T-cell signaling transduction by associating with CD2 and efficiently bringing the Src family protein kinase LCK and LAT to the TCR/CD3 complex (PubMed:19494291). In turn, promotes LCK phosphorylation and subsequent activation (PubMed:12007789). Induces the phosphorylation of the cytoplasmic immunoreceptortyrosine switch motifs (ITSMs) of CD244 initiating a series of signaling events that leads to the generation of the immunological synapse and the directed release of cytolytic granules containing perforin and granzymes by T-lymphocytes and NK- cells (PubMed:27249817).
Cellular Location	Cell membrane; Lipid-anchor, GPI-anchor. Membrane raft. Secreted

## Images



Western blot analysis of CD48 expression in Raji cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.