

## DHRS3 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP9188c

### Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">O75911</a>
<b>Other Accession</b>	<a href="#">O77769</a>
<b>Reactivity</b>	Human, Mouse
<b>Predicted</b>	Bovine
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB23762
<b>Calculated MW</b>	33548
<b>Antigen Region</b>	85-113

### Additional Information

---

<b>Gene ID</b>	9249
<b>Other Names</b>	Short-chain dehydrogenase/reductase 3, DD831, Retinal short-chain dehydrogenase/reductase 1, retSDR1, DHRS3
<b>Target/Specificity</b>	This DHRS3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 85-113 amino acids from the Central region of human DHRS3.
<b>Dilution</b>	WB~~1:1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	DHRS3 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

### Protein Information

---

<b>Name</b>	DHRS3
<b>Synonyms</b>	RDH17, SDR16C1

<b>Function</b>	Catalyzes the reduction of all-trans-retinal to all-trans- retinol in the presence of NADPH.
<b>Cellular Location</b>	Membrane; Multi-pass membrane protein
<b>Tissue Location</b>	Widely expressed with highest levels found in heart, placenta, lung, liver, kidney, pancreas, thyroid, testis, stomach, trachea and spinal cord. Lower levels found in skeletal muscle, intestine and lymph node. No expression detected in brain. In the retina, expressed in cone but not rod outer segments


## Background

DHRS3 catalyze the oxidation/reduction of a wide range of substrates, including retinoids and steroids.


## References

Persson,B., et.al., Chem. Biol. Interact. 178 (1-3), 94-98 (2009)  
Roni,V., et.al., BMC Genomics 8, 42 (2007)

## Images




All lanes : Anti-DHRS3 Antibody (Center) at 1:1000 dilution  
Lane 1: A375 whole cell lysate Lane 2: HepG2 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 34 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



DHRS3 Antibody (Center) (Cat. #AP9188c) western blot analysis in HepG2 cell line and mouse liver tissue lysates (35ug/lane).This demonstrates the DHRS3 antibody detected the DHRS3 protein (arrow).

Western blot analysis of DHRS3 (arrow) using rabbit polyclonal DHRS3 Antibody (Center) (Cat. #AP9188c). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the DHRS3 gene.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.