

Xanthine Oxidase Antibody

Rabbit mAb

Catalog # AP91900

Product Information

Application	WB, IHC
Primary Accession	P47989
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	Xanthine dehydrogenase; Xanthine oxidase; Xanthine oxidoreductase; XD; XDH; xdha; XO; xor;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	146424


Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Xanthine Oxidase
Description	Key enzyme in purine degradation. Catalyzes the oxidation of hypoxanthine to xanthine. Catalyzes the oxidation of xanthine to uric acid. Contributes to the generation of reactive oxygen species. Has also low oxidase activity towards aldehydes (in vitro).
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	XDH
Synonyms	XDHA
Function	Key enzyme in purine degradation. Catalyzes the oxidation of hypoxanthine to xanthine. Catalyzes the oxidation of xanthine to uric acid. Contributes to the generation of reactive oxygen species. Has also low oxidase activity towards aldehydes (in vitro).
Cellular Location	Cytoplasm. Peroxisome. Secreted
Tissue Location	Detected in milk (at protein level). {ECO:0000269 Ref.12}

Images



Western blot analysis of Xanthine Oxidase expression in 293T cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.