

NOXA2/p67phox Antibody

Rabbit mAb

Catalog # AP91907

Product Information

Application	WB, IHC, IF, ICC, IP, IHF
Primary Accession	P19878
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	Ncf2; NOXA2; P67 PHOX;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	59762


Additional Information

Dilution	WB 1:500~1:2000 IHC 1:100~1:500 ICC/IF 1:100~1:500 IP 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human NOXA2/p67phox
Description	NCF2, NCF1, and a membrane bound cytochrome b558 are required for activation of the latent NADPH oxidase (necessary for superoxide production).
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	NCF2 (HGNC:7661)
Function	Subunit of the phagocyte NADPH oxidase complex that mediates the transfer of electrons from cytosolic NADPH to O ₂ to produce the superoxide anion (O ₂ ⁻) (PubMed: 12207919 , PubMed: 38355798). In the activated complex, electrons are first transferred from NADPH to flavin adenine dinucleotide (FAD) and subsequently transferred via two heme molecules to molecular oxygen, producing superoxide through an outer- sphere reaction (PubMed: 38355798). Activation of the NADPH oxidase complex is initiated by the assembly of cytosolic subunits of the NADPH oxidase complex with the core NADPH oxidase complex to form a complex at the plasma membrane or phagosomal membrane (PubMed: 38355798). This activation process is initiated by phosphorylation dependent binding of the cytosolic NCF1/p47-phox subunit to the C-terminus of CYBA/p22-phox (By similarity).
Cellular Location	Cytoplasm.

Images



Western blot analysis of NOXA2/p67phox expression in THP-1 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.