

# LIMK1 Antibody

Rabbit mAb

Catalog # AP92026

## Product Information

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">P53667</a>
<b>Reactivity</b>	Rat, Human, Mouse
<b>Clonality</b>	Monoclonal
<b>Other Names</b>	LIM kinase; LIMK 1; LIMK;
<b>Isotype</b>	Rabbit IgG
<b>Host</b>	Rabbit
<b>Calculated MW</b>	72585

## Additional Information

<b>Dilution</b>	WB 1:500~1:2000
<b>Purification</b>	Affinity-chromatography
<b>Immunogen</b>	A synthesized peptide derived from human LIM Kinase 1
<b>Description</b>	Protein kinase which regulates actin filament dynamics. Phosphorylates and inactivates the actin binding/depolymerizing factor cofilin, thereby stabilizing the actin cytoskeleton. Stimulates axonal outgrowth and may be involved in brain development. Isoform 3 has a dominant negative effect on actin cytoskeletal changes.
<b>Storage Condition and Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

## Protein Information

<b>Name</b>	LIMK1
<b>Synonyms</b>	LIMK
<b>Function</b>	Serine/threonine-protein kinase that plays an essential role in the regulation of actin filament dynamics. Acts downstream of several Rho family GTPase signal transduction pathways (PubMed: <a href="#">10436159</a> , PubMed: <a href="#">11832213</a> , PubMed: <a href="#">12807904</a> , PubMed: <a href="#">15660133</a> , PubMed: <a href="#">16230460</a> , PubMed: <a href="#">18028908</a> , PubMed: <a href="#">22328514</a> , PubMed: <a href="#">23633677</a> ). Activated by upstream kinases including ROCK1, PAK1 and PAK4, which phosphorylate LIMK1 on a threonine residue located in its activation loop (PubMed: <a href="#">10436159</a> ). LIMK1 subsequently phosphorylates and inactivates the actin binding/depolymerizing factors cofilin-1/CFL1, cofilin-2/CFL2 and destrin/DSTN, thereby preventing the cleavage of filamentous actin (F-actin), and stabilizing the actin cytoskeleton (PubMed: <a href="#">11832213</a> , PubMed: <a href="#">15660133</a> , PubMed: <a href="#">16230460</a> , PubMed: <a href="#">23633677</a> ). In this way LIMK1 regulates several

actin-dependent biological processes including cell motility, cell cycle progression, and differentiation (PubMed:[11832213](#), PubMed:[15660133](#), PubMed:[16230460](#), PubMed:[23633677](#)). Phosphorylates TPPP on serine residues, thereby promoting microtubule disassembly (PubMed:[18028908](#)). Stimulates axonal outgrowth and may be involved in brain development (PubMed:[18028908](#)).

#### Cellular Location


Cytoplasm. Nucleus. Cytoplasm, cytoskeleton. Cell projection, lamellipodium {ECO:0000250|UniProtKB:P53668} Note=Predominantly found in the cytoplasm. Localizes in the lamellipodium in a CDC42BPA, CDC42BPB and FAM89B/LRAP25-dependent manner. {ECO:0000250|UniProtKB:P53668}

#### Tissue Location

Highest expression in both adult and fetal nervous system. Detected ubiquitously throughout the different regions of adult brain, with highest levels in the cerebral cortex. Expressed to a lesser extent in heart and skeletal muscle

## Images

---



Western blot analysis of LIM Kinase 1 expression in U-87MG cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.