10320 Camino Santa Fe, Suite G San Diego, CA 92121 Tel: 858.875.1900 Fax: 858.875.1999



# Nck Antibody

Rabbit mAb Catalog # AP92046

### **Product Information**

**Application** WB, IHC, IF, FC, ICC, IP, IHF

**Primary Accession** P16333

Reactivity Rat, Human, Mouse

Clonality Monoclonal

NCK alpha; NCK; NCK1; NCKalpha; **Other Names** 

Isotype Rabbit IgG Host Rabbit **Calculated MW** 42864

## **Additional Information**

**Dilution** WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 IP 1:50 FC 1:500

**Purification** Affinity-chromatography

A synthesized peptide derived from human Nck **Immunogen** 

**Description** Adapter protein which associates with tyrosine-phosphorylated growth factor

receptors or their cellular substrates. Maintains low levels of EIF2S1

phosphorylation by promoting its dephosphorylation by PP1.

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium Storage Condition and Buffer

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

#### **Protein Information**

Name NCK1

**Synonyms** NCK

**Function** Adapter protein which associates with tyrosine-phosphorylated growth

> factor receptors, such as KDR and PDGFRB, or their cellular substrates. Maintains low levels of EIF2S1 phosphorylation by promoting its


dephosphorylation by PP1. Plays a role in the DNA damage response, not in the detection of the damage by ATM/ATR, but for efficient activation of downstream effectors, such as that of CHEK2. Plays a role in ELK1-dependent transcriptional activation in response to activated Ras signaling. Modulates the activation of EIF2AK2/PKR by dsRNA. May play a role in cell adhesion and

migration through interaction with ephrin receptors.

Cytoplasm. Endoplasmic reticulum. Nucleus. Note=Mostly cytoplasmic, but **Cellular Location** 

> shuttles between the cytoplasm and the nucleus. Import into the nucleus requires the interaction with SOCS7 Predominantly nuclear following genotoxic stresses, such as UV irradiation, hydroxyurea or mitomycin C

# **Images**



Western blot analysis of Nck expression in Hela cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.