

KAT1 / HAT1 Antibody

Rabbit mAb

Catalog # AP92128

Product Information

Application	WB, IHC, IF, FC, ICC, IP, IHF
Primary Accession	O14929
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	hat1; KAT1;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	49541

Additional Information

Dilution	WB 1:500~1:1000 IHC 1:50~1:200 ICC/IF 1:50~1:200 IP 1:50 FC 1:100
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human KAT1 / HAT1
Description	Acetylates soluble but not nucleosomal histone H4 at 'Lys-5' (H4K5ac) and 'Lys-12' (H4K12ac) and, to a lesser extent, acetylates histone H2A at 'Lys-5' (H2AK5ac). Has intrinsic substrate specificity that modifies lysine in recognition sequence GXGKXG.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information


Name	HAT1
Synonyms	KAT1
Function	Histone acetyltransferase that plays a role in different biological processes including cell cycle progression, glucose metabolism, histone production or DNA damage repair (PubMed: 20953179 , PubMed: 23653357 , PubMed: 31278053 , PubMed: 32081014). Coordinates histone production and acetylation via H4 promoter binding (PubMed: 31278053). Acetylates histone H4 at 'Lys-5' (H4K5ac) and 'Lys-12' (H4K12ac) and, to a lesser extent, histone H2A at 'Lys-5' (H2AK5ac) (PubMed: 11585814 , PubMed: 22615379). Drives H4 production by chromatin binding to support chromatin replication and acetylation. Since transcription of H4 genes is tightly coupled to S-phase, plays an important role in S-phase entry and progression (PubMed: 31278053). Promotes homologous recombination in DNA repair by facilitating histone turnover and incorporation of acetylated H3.3 at sites of double-strand breaks (PubMed: 23653357). In addition, acetylates other substrates such as

chromatin-related proteins (PubMed:[32081014](#)). Also acetylates RSAD2 which mediates the interaction of ubiquitin ligase UBE4A with RSAD2 leading to RSAD2 ubiquitination and subsequent degradation (PubMed:[31812350](#)).

Cellular Location

[Isoform A]: Nucleus matrix Mitochondrion

Images



Western blot analysis of KAT1 / HAT1 expression in (1) MCF7 cell lysate; (2) NIH/3T3 cell lysate; (3) C6 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.