

IFNGR1 Antibody

Rabbit mAb Catalog # AP92151

Product Information

Application WB, IHC, IF, FC, ICC, IHF

Primary Accession
Reactivity
Human
Clonality
Monoclonal

Other Names CD119; CDw119; IFN gamma R alpha; IFNGR1;

IsotypeRabbit IgGHostRabbitCalculated MW54405

Additional Information

Dilution WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 FC 1:500

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human IFNGR1

Description Receptor for interferon gamma. Two receptors bind one interferon gamma

dimer.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

Protein Information

Name IFNGR1 (HGNC:5439)

Function Receptor subunit for interferon gamma/INFG that plays crucial roles in


antimicrobial, antiviral, and antitumor responses by activating effector immune cells and enhancing antigen presentation (PubMed:20015550). Associates with transmembrane accessory factor IFNGR2 to form a functional

receptor (PubMed: 10986460, PubMed: 2971451, PubMed: 7615558,

PubMed:7617032, PubMed:7673114). Upon ligand binding, the intracellular domain of IFNGR1 opens out to allow association of downstream signaling components JAK1 and JAK2. In turn, activated JAK1 phosphorylates IFNGR1 to form a docking site for STAT1. Subsequent phosphorylation of STAT1 leads to dimerization, translocation to the nucleus, and stimulation of target gene transcription (PubMed:28883123). STAT3 can also be activated in a similar manner although activation seems weaker. IFNGR1 intracellular domain phosphorylation also provides a docking site for SOCS1 that regulates the JAK-STAT pathway by competing with STAT1 binding to IFNGR1 (By similarity).

Cellular Location Cell membrane; Single-pass type I membrane protein

Images



Western blot analysis of IFNGR1 expression in MCF7 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.