

HSF1 Antibody

Rabbit mAb

Catalog # AP92154

Product Information

Application	WB, IHC, IF, FC, ICC, IP, IHF
Primary Accession	Q00613
Reactivity	Human
Clonality	Monoclonal
Other Names	Heat shock factor 1; hsf1; HSTF1;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	57260

Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 IP 1:50 FC 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human HSF1
Description	DNA-binding protein that specifically binds heat shock promoter elements (HSE) and activates transcription. In higher eukaryotes, HSF is unable to bind to the HSE unless the cells are heat shocked.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	HSF1 (HGNC:5224)
Synonyms	HSTF1
Function	Functions as a stress-inducible and DNA-binding transcription factor that plays a central role in the transcriptional activation of the heat shock response (HSR), leading to the expression of a large class of molecular chaperones, heat shock proteins (HSPs), that protect cells from cellular insult damage (PubMed: 11447121 , PubMed: 12659875 , PubMed: 12917326 , PubMed: 15016915 , PubMed: 18451878 , PubMed: 1871105 , PubMed: 1986252 , PubMed: 25963659 , PubMed: 26754925 , PubMed: 7623826 , PubMed: 7760831 , PubMed: 8940068 , PubMed: 8946918 , PubMed: 9121459 , PubMed: 9341107 , PubMed: 9499401 , PubMed: 9535852 , PubMed: 9727490). In unstressed cells, is present in a HSP90-containing multichaperone complex that maintains it in a non-DNA-binding inactivated monomeric form (PubMed: 11583998 , PubMed: 16278218 , PubMed: 9727490). Upon exposure to heat and other stress stimuli, undergoes homotrimerization and activates HSP gene transcription through binding to site-specific heat shock elements (HSEs)

present in the promoter regions of HSP genes (PubMed:[10359787](#), PubMed:[11583998](#), PubMed:[12659875](#), PubMed:[16278218](#), PubMed:[1871105](#), PubMed:[1986252](#), PubMed:[25963659](#), PubMed:[26754925](#), PubMed:[7623826](#), PubMed:[7935471](#), PubMed:[8455624](#), PubMed:[8940068](#), PubMed:[9499401](#), PubMed:[9727490](#)). Upon heat shock stress, forms a chromatin-associated complex with TTC5/STRAP and p300/EP300 to stimulate HSR transcription, therefore increasing cell survival (PubMed:[18451878](#)). Activation is reversible, and during the attenuation and recovery phase period of the HSR, returns to its unactivated form (PubMed:[11583998](#), PubMed:[16278218](#)). Binds to inverted 5'-NGAAN-3' pentamer DNA sequences (PubMed:[1986252](#), PubMed:[26727489](#)). Binds to chromatin at heat shock gene promoters (PubMed:[25963659](#)). Activates transcription of transcription factor FOXR1 which in turn activates transcription of the heat shock chaperones HSPA1A and HSPA6 and the antioxidant NADPH-dependent reductase DHRS2 (PubMed:[34723967](#)). Also serves several other functions independently of its transcriptional activity. Involved in the repression of Ras-induced transcriptional activation of the c-fos gene in heat-stressed cells (PubMed:[9341107](#)). Positively regulates pre-mRNA 3'-end processing and polyadenylation of HSP70 mRNA upon heat-stressed cells in a symplekin (SYMPK)-dependent manner (PubMed:[14707147](#)). Plays a role in nuclear export of stress-induced HSP70 mRNA (PubMed:[17897941](#)). Plays a role in the regulation of mitotic progression (PubMed:[18794143](#)). Also plays a role as a negative regulator of non-homologous end joining (NHEJ) repair activity in a DNA damage-dependent manner (PubMed:[26359349](#)). Involved in stress-induced cancer cell proliferation in a IER5-dependent manner (PubMed:[26754925](#)).


Cellular Location

Nucleus. Cytoplasm. Nucleus, nucleoplasm. Cytoplasm, perinuclear region. Cytoplasm, cytoskeleton, spindle pole. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Chromosome, centromere, kinetochore
Note=The monomeric form is cytoplasmic in unstressed cells (PubMed:[26159920](#), PubMed:[8455624](#)). Predominantly nuclear protein in both unstressed and heat shocked cells (PubMed:[10359787](#), PubMed:[10413683](#)). Translocates in the nucleus upon heat shock (PubMed:[8455624](#)). Nucleocytoplasmic shuttling protein (PubMed:[26159920](#)). Colocalizes with IER5 in the nucleus (PubMed:[27354066](#)). Colocalizes with BAG3 to the nucleus upon heat stress (PubMed:[26159920](#), PubMed:[8455624](#)). Localizes in subnuclear granules called nuclear stress bodies (nSBs) upon heat shock (PubMed:[10359787](#), PubMed:[10747973](#), PubMed:[11447121](#), PubMed:[11514557](#), PubMed:[19229036](#), PubMed:[24581496](#), PubMed:[25963659](#)). Colocalizes with SYMPK and SUMO1 in nSBs upon heat shock (PubMed:[10359787](#), PubMed:[11447121](#), PubMed:[11514557](#), PubMed:[12665592](#), PubMed:[14707147](#)) Colocalizes with PRKACA/PKA in the nucleus and nSBs upon heat shock (PubMed:[21085490](#)). Relocalizes from the nucleus to the cytoplasm during the attenuation and recovery phase period of the heat shock response (PubMed:[26159920](#)). Translocates in the cytoplasm in a YWHAE- and XPO1/CRM1-dependent manner (PubMed:[12917326](#)). Together with histone H2AX, redistributed in discrete nuclear DNA damage-induced foci after ionizing radiation (IR) (PubMed:[26359349](#)). Colocalizes with calcium-responsive transactivator SS18L1 at kinetochore region on the mitotic chromosomes (PubMed:[18794143](#)). Colocalizes with gamma tubulin at centrosome (PubMed:[18794143](#)). Localizes at spindle pole in metaphase (PubMed:[18794143](#)). Colocalizes with PLK1 at spindle poles during prometaphase (PubMed:[18794143](#)).

Images

Western blot analysis of HSF1 expression in HeLa cell

lysate.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.