

SQSTM1 / p62 Antibody

Rabbit mAb

Catalog # AP92155

Product Information

Application	WB, IHC, IF, FC, ICC, IP, IHF
Primary Accession	Q13501
Reactivity	Human
Clonality	Monoclonal
Other Names	A170; DMRV; EBIAP; ORCA; OSF-6; Osi; OSIL; p60; p62; p62B; PDB3; Sequestosome 1; Sqstm1; STAP; STONE14; Ubiquitin binding protein p62; ZIP; ZIP3;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	47687

Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 IP 1:50 FC 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human SQSTM1 / p62
Description	Adapter protein which binds ubiquitin and may regulate the activation of NFkB1 by TNF-alpha, nerve growth factor (NGF) and interleukin-1. May play a role in titin/TTN downstream signaling in muscle cells. May regulate signaling cascades through ubiquitination. Adapter that mediates the interaction between TRAF6 and CYLD (By similarity).
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	SQSTM1 {ECO:0000303 PubMed:16286508, ECO:0000312 HGNC:HGNC:11280}
Function	Molecular adapter required for selective macroautophagy (aggrephagy) by acting as a bridge between polyubiquitinated proteins and autophagosomes (PubMed: 15340068 , PubMed: 15953362 , PubMed: 16286508 , PubMed: 17580304 , PubMed: 20168092 , PubMed: 22017874 , PubMed: 22622177 , PubMed: 24128730 , PubMed: 28404643 , PubMed: 29343546 , PubMed: 29507397 , PubMed: 31857589 , PubMed: 33509017 , PubMed: 34471133 , PubMed: 34893540 , PubMed: 35831301 , PubMed: 37306101 , PubMed: 37802024). Promotes the recruitment of ubiquitinated cargo proteins to autophagosomes via multiple domains that bridge proteins and organelles in different steps (PubMed: 16286508 , PubMed: 20168092 , PubMed: 22622177 ,

PubMed:[24128730](#), PubMed:[28404643](#), PubMed:[29343546](#), PubMed:[29507397](#), PubMed:[34893540](#), PubMed:[37802024](#)). SQSTM1 first mediates the assembly and removal of ubiquitinated proteins by undergoing liquid-liquid phase separation upon binding to ubiquitinated proteins via its UBA domain, leading to the formation of insoluble cytoplasmic inclusions, known as p62 bodies (PubMed:[15911346](#), PubMed:[20168092](#), PubMed:[22017874](#), PubMed:[24128730](#), PubMed:[29343546](#), PubMed:[29507397](#), PubMed:[31857589](#), PubMed:[37802024](#)). SQSTM1 then interacts with ATG8 family proteins on autophagosomes via its LIR motif, leading to p62 body recruitment to autophagosomes, followed by autophagic clearance of ubiquitinated proteins (PubMed:[16286508](#), PubMed:[17580304](#), PubMed:[20168092](#), PubMed:[22622177](#), PubMed:[24128730](#), PubMed:[28404643](#), PubMed:[37802024](#)). SQSTM1 is itself degraded along with its ubiquitinated cargos (PubMed:[16286508](#), PubMed:[17580304](#), PubMed:[37802024](#)). Also required to recruit ubiquitinated proteins to PML bodies in the nucleus (PubMed:[20168092](#)). Also involved in autophagy of peroxisomes (pexophagy) in response to reactive oxygen species (ROS) by acting as a bridge between ubiquitinated PEX5 receptor and autophagosomes (PubMed:[26344566](#)). Acts as an activator of the NFE2L2/NRF2 pathway via interaction with KEAP1: interaction inactivates the BCR(KEAP1) complex by sequestering the complex in inclusion bodies, promoting nuclear accumulation of NFE2L2/NRF2 and subsequent expression of cytoprotective genes (PubMed:[20452972](#), PubMed:[28380357](#), PubMed:[33393215](#), PubMed:[37306101](#)). Promotes relocalization of 'Lys-63'-linked ubiquitinated STING1 to autophagosomes (PubMed:[29496741](#)). Involved in endosome organization by retaining vesicles in the perinuclear cloud: following ubiquitination by RNF26, attracts specific vesicle-associated adapters, forming a molecular bridge that restrains cognate vesicles in the perinuclear region and organizes the endosomal pathway for efficient cargo transport (PubMed:[27368102](#), PubMed:[33472082](#)). Sequesters tensin TNS2 into cytoplasmic puncta, promoting TNS2 ubiquitination and proteasomal degradation (PubMed:[25101860](#)). May regulate the activation of NFkB1 by TNF-alpha, nerve growth factor (NGF) and interleukin-1 (PubMed:[10356400](#), PubMed:[10747026](#), PubMed:[11244088](#), PubMed:[12471037](#), PubMed:[16079148](#), PubMed:[19931284](#)). May play a role in titin/TTN downstream signaling in muscle cells (PubMed:[15802564](#)). Adapter that mediates the interaction between TRAF6 and CYLD (By similarity).

Cellular Location


Cytoplasmic vesicle, autophagosome. Preautophagosomal structure. Cytoplasm, cytosol. Nucleus, PML body. Late endosome. Lysosome. Nucleus Endoplasmic reticulum. Cytoplasm, myofibril, sarcomere {ECO:0000250|UniProtKB:O08623}. Note=In cardiac muscle, localizes to the sarcomeric band (By similarity). Localizes to cytoplasmic membraneless inclusion bodies, known as p62 bodies, containing polyubiquitinated protein aggregates (PubMed:[11786419](#), PubMed:[20357094](#), PubMed:[22017874](#), PubMed:[29343546](#), PubMed:[29507397](#), PubMed:[31857589](#), PubMed:[37306101](#), PubMed:[37802024](#)). In neurodegenerative diseases, detected in Lewy bodies in Parkinson disease, neurofibrillary tangles in Alzheimer disease, and HTT aggregates in Huntington disease (PubMed:[15158159](#)). In protein aggregate diseases of the liver, found in large amounts in Mallory bodies of alcoholic and nonalcoholic steatohepatitis, hyaline bodies in hepatocellular carcinoma, and in SERPINA1 aggregates (PubMed:[11981755](#)) Enriched in Rosenthal fibers of pilocytic astrocytoma (PubMed:[11786419](#)). In the cytoplasm, observed in both membrane-free ubiquitin-containing protein aggregates (sequestosomes) and membrane-surrounded autophagosomes (PubMed:[15953362](#), PubMed:[17580304](#)) Colocalizes with TRIM13 in the perinuclear endoplasmic reticulum (PubMed:[22178386](#)). Co-localizes with TRIM5 in cytoplasmic bodies (PubMed:[20357094](#)). When nuclear export is blocked by treatment with leptomycin B, accumulates in PML bodies (PubMed:[20168092](#))

{ECO:0000250|UniProtKB:O08623, ECO:0000269|PubMed:11786419,
ECO:0000269|PubMed:11981755, ECO:0000269|PubMed:15158159,
ECO:0000269|PubMed:15953362, ECO:0000269|PubMed:17580304,
ECO:0000269|PubMed:20168092, ECO:0000269|PubMed:20357094,
ECO:0000269|PubMed:22017874, ECO:0000269|PubMed:22178386,
ECO:0000269|PubMed:29343546, ECO:0000269|PubMed:29507397,
ECO:0000269|PubMed:31857589, ECO:0000269|PubMed:37306101,
ECO:0000269|PubMed:37802024}

Tissue Location

Ubiquitously expressed.

Images



Western blot analysis of SQSTM1 / p62 expression in HepG2 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.