

# RAIDD Antibody

Rabbit mAb

Catalog # AP92160

## Product Information

---

|                          |                        |
|--------------------------|------------------------|
| <b>Application</b>       | WB, IF, FC, ICC, IP    |
| <b>Primary Accession</b> | <a href="#">P78560</a> |
| <b>Reactivity</b>        | Rat, Human             |
| <b>Clonality</b>         | Monoclonal             |
| <b>Other Names</b>       | Cradd;                 |
| <b>Isotype</b>           | Rabbit IgG             |
| <b>Host</b>              | Rabbit                 |
| <b>Calculated MW</b>     | 22745                  |

## Additional Information

---

|                                     |   |
|-------------------------------------|---|
| <b>Dilution</b>                     | WB 1:500~1:2000 ICC/IF 1:50~1:200 IP 1:50 FC 1:50   |
| <b>Purification</b>                 | Affinity-chromatography   |
| <b>Immunogen</b>                    | A synthesized peptide derived from human RAIDD  |
| <b>Description</b>                  | Apoptotic adaptor molecule specific for caspase-2 and FASL/TNF receptor-interacting protein RIP. In the presence of RIP and TRADD, CRADD recruits caspase-2 to the TNFR-1 signalling complex. |
| <b>Storage Condition and Buffer</b> | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.             |


## Protein Information

---

|                          |  |
|--------------------------|--|
| <b>Name</b>              | CRADD  |
| <b>Synonyms</b>          | RAIDD  |
| <b>Function</b>          | Adapter protein that associates with PIDD1 and the caspase CASP2 to form the PIDDosome, a complex that activates CASP2 and triggers apoptosis (PubMed: <a href="#">15073321</a> , PubMed: <a href="#">16652156</a> , PubMed: <a href="#">17159900</a> , PubMed: <a href="#">17289572</a> , PubMed: <a href="#">9044836</a> ). Also recruits CASP2 to the TNFR-1 signaling complex through its interaction with RIPK1 and TRADD and may play a role in the tumor necrosis factor-mediated signaling pathway (PubMed: <a href="#">8985253</a> ). |
| <b>Cellular Location</b> | Cytoplasm {ECO:0000250 UniProtKB:O88843}. Nucleus {ECO:0000250 UniProtKB:O88843}   |
| <b>Tissue Location</b>   | Constitutively expressed in most tissues, with particularly high expression in adult heart, testis, liver, skeletal muscle, fetal liver and kidney.  |

## Images

---



Western blot analysis of RAIDD expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.