

TPPP Antibody

Rabbit mAb

Catalog # AP92165

Product Information

Application	WB, IHC, IF, FC, ICC, IHF
Primary Accession	O94811
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Other Names	p24; p25; p25alpha; TPPP; TPPP/p25; TPPP1; Tubulin polymerization promoting protein;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	23694

Additional Information

Dilution	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 FC 1:50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human TPPP
Description	May play a role in the polymerization of tubulin into microtubules, microtubule bundling and the stabilization of existing microtubules, thus maintaining the integrity of the microtubule network. May play a role in mitotic spindle assembly and nuclear envelope breakdown.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Protein Information

Name	TPPP {ECO:0000303 PubMed:17105200, ECO:0000312 HGNC:HGNC:24164}
Function	Regulator of microtubule dynamics that plays a key role in myelination by promoting elongation of the myelin sheath (PubMed: 31522887). Acts as a microtubule nucleation factor in oligodendrocytes: specifically localizes to the postsynaptic Golgi apparatus region, also named Golgi outpost, and promotes microtubule nucleation, an important step for elongation of the myelin sheath (PubMed: 31522887 , PubMed: 33831707). Required for both uniform polarized growth of distal microtubules as well as directing the branching of proximal processes (PubMed: 31522887). Shows magnesium-dependent GTPase activity; the role of the GTPase activity is unclear (PubMed: 21316364 , PubMed: 21995432). In addition to microtubule nucleation activity, also involved in microtubule bundling and stabilization of existing microtubules, thereby maintaining the integrity of the microtubule network (PubMed: 17105200 , PubMed: 17693641 , PubMed: 18028908 , PubMed: 26289831). Regulates microtubule dynamics by promoting tubulin

acetylation: acts by inhibiting the tubulin deacetylase activity of HDAC6 (PubMed:[20308065](#), PubMed:[23093407](#)). Also regulates cell migration: phosphorylation by ROCK1 inhibits interaction with HDAC6, resulting in decreased acetylation of tubulin and increased cell motility (PubMed:[23093407](#)). Plays a role in cell proliferation by regulating the G1/S-phase transition (PubMed:[23355470](#)). Involved in astral microtubule organization and mitotic spindle orientation during early stage of mitosis; this process is regulated by phosphorylation by LIMK2 (PubMed:[22328514](#)).


Cellular Location

Golgi outpost {ECO:0000250|UniProtKB:D3ZQL7}. Cytoplasm, cytoskeleton, microtubule organizing center {ECO:0000250|UniProtKB:D3ZQL7}. Cytoplasm, cytoskeleton. Nucleus Cytoplasm, cytoskeleton, spindle Note=Specifically localizes to the postsynaptic Golgi apparatus region, also named Golgi outpost, which shapes dendrite morphology by functioning as sites of acentrosomal microtubule nucleation (By similarity). Mainly localizes to the cytoskeleton (PubMed:18028908) Also found in the nucleus; however, nuclear localization is unclear and requires additional evidences (PubMed:18028908). Localizes to glial Lewy bodies in the brains of individuals with synucleinopathies (PubMed:15590652, PubMed:17027006). During mitosis, colocalizes with LIMK2 at the mitotic spindle (PubMed:22328514) {ECO:0000250|UniProtKB:D3ZQL7, ECO:0000269|PubMed:15590652, ECO:0000269|PubMed:17027006, ECO:0000269|PubMed:18028908, ECO:0000269|PubMed:22328514}

Tissue Location

Widely expressed..

Images



Western blot analysis of TPPP expression in SHSY5Y cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.