

## **BOB1** Antibody

Rabbit mAb Catalog # AP92177

## **Product Information**

Application	WB, IHC, IF, FC, ICC, IHF
Primary Accession	<u>Q16633</u>
Reactivity	Human
Clonality	Monoclonal
Other Names	BOB 1; OBF1; OCAB; OCT binding factor 1; Pou2af1;
lsotype	Rabbit IgG
Host	Rabbit
Calculated MW	27436

## **Additional Information**

Dilution Purification	WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 FC 1:50 Affinity-chromatography
Immunogen	A synthesized peptide derived from human BOB1
Description	Transcriptional coactivator that specifically associates with either OCT1 or
Storage Condition and Buffer	OCT2. It boosts the OCT1 mediated promoter activity and to a lesser extent, that of OCT2. It has no intrinsic DNA-binding activity. Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.
	Avoid freeze / thaw cycle.

## **Protein Information**

Name	POU2AF1 ( <u>HGNC:9211</u> )
Function	Transcriptional coactivator that specifically associates with either POU2F1/OCT1 or POU2F2/OCT2 (PubMed: <u>7859290</u> ). It boosts the POU2F1/OCT1 mediated promoter activity and to a lesser extent, that of POU2F2/OCT2 (PubMed: <u>7779176</u> ). It recognizes the POU domains of POU2F1/OCT1 and POU2F2/OCT2 (PubMed: <u>7779176</u> ). It is essential for the response of B-cells to antigens and required for the formation of germinal centers (PubMed: <u>7623806</u> , PubMed: <u>7859290</u> ). Regulates IL6 expression in B cells as POU2F2/OCT2 coactivator (By similarity).
Cellular Location	Nucleus.
Tissue Location	B-cell specific (PubMed:7779176, PubMed:7859290). Detected in mainly in spleen, but also in thymus, periphral blood leukocyte and small intestine (PubMed:7779176, PubMed:7859290)



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.