

ENTPD5 Antibody

Rabbit mAb Catalog # AP92195

Product Information

Application WB, IHC, IP **Primary Accession** 075356

Reactivity Rat, Human, Mouse

Clonality Monoclonal

Other Names CD39L4; Entpd5; mNTPase; NTPDase 5; PCPH; UDPase ENTPD5;

IsotypeRabbit IgGHostRabbitCalculated MW47517

Additional Information

Dilution WB 1:500~1:2000 IHC 1:50~1:200 IP 1:50

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human ENTPD5

Description Uridine diphosphatase (UDPase) that promotes protein N-glycosylation and

ATP level regulation. UDP hydrolysis promotes protein N-glycosylation and folding in the endoplasmic reticulum, as well as elevated ATP consumption in

the cytosol via an ATP hydrolysis cycle.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

Protein Information

Name ENTPD5 (HGNC:3367)

Function Hydrolyzes nucleoside diphosphates with a preference for GDP, IDP and

UDP compared to ADP and CDP (PubMed: <u>10400613</u>, PubMed: <u>15698960</u>). In the lumen of the endoplasmic reticulum, hydrolyzes UDP that acts as an

end-product feedback inhibitor of the UDP-Glc:glycoprotein

glucosyltransferases. UMP can be transported back by an UDP-sugar antiporter to the cytosol where it is consumed to regenerate UDP- glucose. Therefore, it positively regulates protein reglucosylation by clearing UDP from the ER lumen and by promoting the regeneration of UDP-glucose. Protein reglucosylation is essential to proper glycoprotein folding and quality control


in the ER (By similarity).

Cellular Location Endoplasmic reticulum {ECO:0000250 | UniProtKB:Q9WUZ9}. Secreted

Tissue Location Expressed in adult liver, kidney, prostate, testis and colon. Much weaker

expression in other tissues

Images



Western blot analysis of ENTPD5 expression in fetal liver lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.