

Oct-2 Antibody

Rabbit mAb Catalog # AP92208

Product Information

Application WB, IHC, IF, FC, ICC, IHF

Primary Accession P09086
Reactivity Human
Clonality Monoclonal

Other Names Oct 2; Octamer binding transcription factor 2; OTF2; POU domain; POU2F2;

IsotypeRabbit IgGHostRabbitCalculated MW51209

Additional Information

Dilution WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200 FC 1:50

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human Oct-2

Description Transcription factor that specifically binds to the octamer motif

(5'-ATTTGCAT-3'). Regulates transcription in a number of tissues in addition to

activating immunoglobulin gene expression. Modulates transcription

transactivation by NR3C1, AR and PGR.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium

azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term.

Avoid freeze / thaw cycle.

Protein Information

Name POU2F2 (HGNC:9213)

Synonyms OCT2, OTF2

Function Transcription factor that specifically binds to the octamer motif

(5'-ATTTGCAT-3') (PubMed:2904654, PubMed:7859290). Regulates IL6

expression in B cells with POU2AF1 (By similarity). Regulates transcription in a number of tissues in addition to activating immunoglobulin gene expression (PubMed:2901913, PubMed:2904654). Modulates transcription transactivation

by NR3C1, AR and PGR (PubMed: 10480874).


Cellular Location Cytoplasm. Nucleus.

Tissue Location Isoform 3 is B-cell specific. Isoform 5 is expressed in B-cells and the

immunoglobulin-expressing T-cell line MOLT-4, but not in the T-cell line

BW5147.

Images



Western blot analysis of Oct-2 expression in Daudi cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.