10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# TNFSF9 Antibody

Rabbit mAb
Catalog # AP92219

## Product Information

| | |
|---|---|
| **Application** | WB, IP |
| **Primary Accession** | P41273 |
| **Reactivity** | Human |
| **Clonality** | Monoclonal |
| **Other Names** | 4 1BB L; 4 1BBL; 4-1BB ligand; 4-1BBL; Cd137l; Cd157l; Ly63l; Tnfsf9; TNLG5A; |
| **Isotype** | Rabbit IgG |
| **Host** | Rabbit |
| **Calculated MW** | 26625 |

## Additional Information


| | |
|---|---|
| **Dilution** | WB 1:500~1:2000 IP 1:50 |
| **Purification** | Affinity-chromatography |
| **Immunogen** | A synthesized peptide derived from human TNFSF9 |
| **Description** | Cytokine that binds to TNFRSF9. Induces the proliferation of activated peripheral blood T-cells. May have a role in activation-induced cell death (AICD). May play a role in cognate interactions between T-cells and B-cells/macrophages. |
| **Storage Condition and Buffer** | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |

## Protein Information

| | |
|---|---|
| **Name** | TNFSF9 |
| **Function** | Cytokine that binds to TNFRSF9. Induces the proliferation of activated peripheral blood T-cells. May have a role in activation- induced cell death (AICD). May play a role in cognate interactions between T-cells and B-cells/macrophages. |
| **Cellular Location** | Membrane; Single-pass type II membrane protein. |
| **Tissue Location** | Expressed in brain, placenta, lung, skeletal muscle and kidney |

## Images

Western blot analysis of TNFSF9 expression in HEK293 cell lysate.




Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.